# Existence and uniqueness of H-systems's solutions with Dirichlet conditions

**P. Amster, M. C. Mariani and D. F. Rial**

ABSTRACT: *We give conditions on the boundary data in order to obtain a solution for the problem (1) below and we prove that in certain cases the solutions are isolated. The condition* $\|H\|_\infty \|g\|_\infty \leq 1$ *is not necessary.*

## Introduction

We consider the Dirichlet problem in the unit disc $B = \{(u,v) \in R^2; u^2+v^2 < 1\}$ for a vector function $X : \overline{B} \longrightarrow R^3$ which satisfies the equation of prescribed mean curvature

$$(1) \begin{cases} \Delta X = 2H(u,v) X_u \wedge X_v & \text{in} \quad B \\ X = g & \text{in} \quad \partial B \end{cases}$$

where $X_u = \dfrac{\partial X}{\partial u}$, $X_v = \dfrac{\partial X}{\partial v}$, $\wedge$ denotes the exterior product in $R^3$ and $H : \overline{B} \longrightarrow R$ is a given continuous function.

The problem above arises in the Plateau and Dirichlet problems for the prescribed mean curvature equation that has been studied in [1-3-4-5-6-7-8].

The main results are the following theorems:

THEOREM 1 *Let be* $2 < p < \infty$, *and* $g \in W^{2,p}(B, R^3)$ *such that* $g - a$ *is small enough for a certain* $a \in R$, *then there exists a solution* $X \in W^{2,p}$ *of (1).*

THEOREM 2 *Let* $X$ *be a solution of (1) in* $W^{2,p}$, $1 < p < \infty$. *Then:*

*i) If* $p \leq 2$ *and* $X \in W^{1,\infty}$ *then* $X$ *is isolated in* $(W^{1,\infty} \cap W^{2,p}, \|\cdot\|_{1,\infty})$.

*ii) If* $p > 2$ *then* $X$ *is isolated in* $W^{1,p}$. *Moreover, if* $\overline{\Omega}$ *is a bounded subset of* $W^{1,p}$ *the number of solutions of (1) in* $\overline{\Omega}$ *is finite.*

REMARK

*For $H$ constant Hildebrandt found a solution of (1) in $W^{1,2}$, for the case $|H|\|g\|_\infty \leq 1$ [3]. Theorem 1 gives a solution for $p > 2$ when $g$ is close to a constant $a$. Since the equation in $B$ depends only on the derivatives of $X$, we may suppose $a = 0$*

## 1. Solution by fixed point methods

The systems (2) and (3) are equivalent to (1) with $X = X_0 + Y$

$$(2)\quad \begin{cases} \Delta X_0 = 0 \text{ in} \quad B \\ X_0 = g \qquad \text{in} \quad \partial B \end{cases}$$

$$(3)\quad \begin{cases} \Delta Y = F(X_0, Y) \qquad \text{in} \quad B \\ Y = 0 \qquad \text{in} \quad \partial B \end{cases}$$

where $F$ is given by

$$F(X_0, Y) = 2H(u,v)[(X_{0u} \wedge Y_v + Y_u \wedge X_{0v}) + (Y_u \wedge Y_v + X_{0u} \wedge X_{0v})]$$

For $g \in W^{2,p}\left(B, R^3\right)$, (2) admits a unique solution in $W^{2,p}\left(B, R^3\right)$. We can rewrite (3) as:

$$\begin{cases} L(X_0)Y = F_1(X_0, Y) \qquad \text{in} \quad B \\ Y = 0 \qquad \text{in} \quad \partial B \end{cases}$$

where $L(X_0)$ is the linear operator

$$L(X_0)Y = \Delta Y - 2H(u,v)(X_{0u} \wedge Y_v + Y_u \wedge X_{0v})$$

and $F_1$ is defined by

$$F_1(X_0, Y) = 2H(u,v)(Y_u \wedge Y_v + X_{0u} \wedge X_{0v}).$$

REMARK $L(X_0)$ *is strictly elliptic and for $p > 2$ its coefficients are bounded since $X_0 \in W^{1,\infty}\left(B, R^3\right)$ and $H \in C(\overline{B})$.*

*We will use the following technical lemmas:*

LEMMA 4 *Let be* $X_0 \in W^{2,p}(B, R^3)$, *then*

*i) If* $2 < p < \infty$,

$$\|F_1(X_0, Y_1)\|_{p/2} \leq 2\|H\|_\infty(\|X_0\|^2_{1,p} + \|Y_1\|^2_{1,p}) \tag{4}$$

$$\|F_1(X_0, Y_1) - F_1(X_0, Y_2)\|_{p/2} \leq 4\|H\|_\infty R\left(\|Y_1 - Y_2\|_{1,p}\right) \tag{5}$$

*for* $Y_1, Y_2 \in B_R(0) \subset W^{1,p}$.

*ii) If* $p > 1$, $X_0$, $Y_1$, $Y_2 \in W^{1,p} \cap W^{1,\infty}$

$$\|F_1(X_0, Y_1)\|_p \leq 2\|H\|_\infty(\|X_0\|_{1,\infty}.\|X_0\|_{1,p} + \|Y_1\|_{1,\infty}.\|Y_1\|_{1,p}) \tag{6}$$

$$\|F_1(X_0, Y_1) - F_1(X_0, Y_2)\|_p \leq 4\|H\|_\infty R\left(\|Y_1 - Y_2\|_{1,p}\right) \tag{7}$$

*for* $Y_1, Y_2 \in B_R(0) \subset W^{1,\infty}$.

*Proof*

$$\|F_1(X_0, Y_1)\|_{p/2} \leq 2\|H\|_\infty(\|Y_{1u} \wedge Y_{1v}\|_{p/2} + \|X_{0u} \wedge X_{0v}\|_{p/2})$$

and

$$\|Y_{1u} \wedge Y_{1v}\|^{p/2}_{p/2} \leq \int |Y_{1u}|^{p/2}|Y_{1v}|^{p/2} \leq \|Y_{1u}\|^{p/2}_p\|Y_{1v}\|^{p/2}_p$$

The same inequality holds for $X_0$, and in order to prove (5), we also have that

$$\|Y_{1u} \wedge Y_{1v} - Y_{2u} \wedge Y_{2v}\|_{p/2} \leq \|Y_{1u} \wedge (Y_{1v} - Y_{2v})\|_{p/2} + \|Y_{2v} \wedge (Y_{1u} - Y_{2u})\|_{p/2}$$

and the proof follows.

In the same way we obtain (6) and (7), using that $\|a \wedge b\|_p \leq \|a\|_\infty.\|b\|_p$ ♣

LEMMA 5 *If* $\|X_0\|_{1,p} \leq \delta$, $2 < p < \infty$ *with* $\delta$ *small enough, then there exists a constant* $C$ *independent of* $X_0$ *verifying*

$$\|Y\|_{2,p/2} \leq C\|L(X_0)Y\|_{p/2}$$

*for all* $Y \in W^{2,p/2} \cap W_0^{1,p/2}$

*Proof*

Writing $L(X_0)Y = \Delta Y - R(X_0)Y$ it follows that

$$\|L(X_0)Y\|_{p/2} \geq \|\Delta Y\|_{p/2} - \|R(X_0)Y\|_{p/2} \geq \frac{1}{C_1}\|Y\|_{2,p/2} - \|R(X_0)Y\|_{p/2}$$

where $C_1$ is the constant provided by Lemma 9.17 in [2].

As $\|R(X_0)Y\|_{p/2} \leq 4\|H\|_\infty \|X_0\|_{1,p}\|Y\|_{1,p} \leq 4\overline{c}\|H\|_\infty \|X_0\|_{1,p}\|Y\|_{2,p/2}$ (where $\overline{c}$ is the constant of Sobolev inmersion $W^{2,p/2} \hookrightarrow W^{1,p}$), the proof follows. ♣

REMARK

*The same Lemma 9.17 in [2] shows that if* $\|g\|_{2,p}$ *is small, then* $\|X_0 - g\|_{2,p}$ *is small, and then also* $\|X_0\|_{1,p}$.

PROPOSITION 6 *Let be* $X_0 \in W^{2,p}\left(B, R^3\right)$ *with* $p > 2$, *then the following problem*

$$(8) \quad \begin{cases} L(X_0)Y = F_1(X_0, \overline{Y}) & in \quad B \\ Y = 0 \qquad in \quad \partial B \end{cases}$$

*define a continuous map* $T : \overline{Y} \longrightarrow Y$ *in* $W_0^{1,p}$. *Furthermore, if* $\|X_0\|_{1,p}$ *is small enough, there exists a number* $R$ *such that* $T$ *is a contraction in* $B_R(0) \subset W^{1,p}$.

*Proof*

¿From Theorem 9.15 in [2], problem (8) admits a unique solution $Y \in W^{2,p/2}$, so the map $T$ is well defined. Also we have from (5) that if $\overline{Y}, \overline{Z} \in B_R \subset W^{1,p}$ then

$$\|Y - Z\|_{2,p/2} \leq C(\|L(X_0)(Y-Z)\|_{p/2} = C\|F_1(X_0, \overline{Y}) - F_1(X_0, \overline{Z})\|_{p/2} \leq 4C\|H\|_\infty R\|\overline{Y} - \overline{Z}\|_{1,p}$$

and using the Sobolev inmersion of $W^{2,p/2}$ in $W^{1,p}$ the continuity of $T$ follows.

Also we have from (4)

$$\|Y\|_{1,p} \leq \overline{c}\,\|Y\|_{2,p/2} \leq C\overline{c}\,\|L(X_0)Y\|_{p/2} \leq 2C\overline{c}\|H\|_\infty(\|X_0\|_{1,p}^2 + \left\|\overline{Y}\right\|_{1,p}^2)$$

Choosing $\|X_0\|_{1,p}$ and $R$ small enough, we obtain

$$4C\|H\|_\infty R < 1$$

$$\|Y\|_{1,p} \leq R$$

and $T$ is a contraction in $B_R$. ♣

REMARK

*Note that if $Y$ is a fixed point of $T$, then $Y \in W^{2,p}$, and we obtain Theorem 1 as an immediate consequence of Proposition 6.*

## 2. Local unicity of the solutions

Now we will prove theorem 2.

Let $X_0$ be a solution of (1). If $Y$ is another solution of (1) then $Z = Y - X_0$ satisfies

$$\begin{cases} L(X_0)Z = 2H(u,v)Z_u \wedge Z_v \text{ in} & B \\ Z = 0 \qquad \text{in} & \partial B \end{cases}$$

We consider the associated problem

$$\begin{cases} L(X_0)Z = 2H(u,v)\overline{Z}_u \wedge \overline{Z}_v \text{ in} & B \\ Z = 0 \qquad \text{in} & \partial B \end{cases}$$

For ii), in the same way as before, we obtain

$$\|Z\|_{1,p} \leq 2C(X_0)\overline{c}\|H\|_\infty \|\overline{Z}\|_{1,p}^2$$

and if $\overline{Z}_1, \overline{Z}_2 \in B_R(0) \subset W^{1,p}$ then

$$\|Z_1 - Z_2\|_{1,p} \leq 4C(X_0)\overline{c}\|H\|_\infty R\|\overline{Z}_1 - \overline{Z}_2\|_{1,p}$$

If we choose $R$ verifying

$$4C(X_0)\overline{c}\|H\|_\infty R < 1$$

the map $\overline{T} : \overline{Z} \longrightarrow Z$ is a contraction in $B_R$, so $X_0$ is isolated.

REMARK *¿From Sobolev inmersions $\overline{T}$ is compact in $W^{1,p}$ and being any solution of (1) a fixed point of $\overline{T}$, we conclude that the number of solutions in $B_R$ is finite for every $R$.*

*In the same way we prove i) using (6) and (7).* ♣

REMARK *The number $R$ in the theorems above can be estimated in terms of the Sobolev inmersions constants, $\|H\|_\infty$ and $\|g\|_{2,p}$.*

**References**


[1] H. Brezis, J. Coron, Multiple solutions of $H$ systems and Rellich's conjecture, Comm. Pure Appl. Math. 37 (1984), 149-187.

[2] D. Gilbarg, N.S. Trudinger, Elliptic partial differential equations of second order, Springer- Verlag (1983).

[3] S. Hildebrandt, On the Plateau problem for surfaces of constant mean curvature. Comm. Pure Appl. Math. 23 (1970) 97-114.

[4] E. Lami Dozo, M.C. Mariani, A Dirichlet problem for an $H$ system with variable $H$. Manuscripta Math. 81 (1993), 1-14.

[5] M. C. Mariani, D. Rial, Solutions to the mean cuvature equation by fixed point methods. Bulletin of The Belgian Mathematical Society - Simon Stevin (1997).

[6] M. Struwe, Plateau 's problem and the calculus of variations, Lecture Notes Princeton Univ. Press (1988).

[7] M. Struwe, Multiple solutions to the Dirichlet problem for the equation of prescribed mean curvature, Research papers published in honor of J. Moser's 60 Birthday, Rabinowitz, P and Ehnder, E. (Eds.), Boston, Academic Press, 639-666 (1990).

[8] Wang Guofang: The Dirichlet problem for the equation of prescribed mean curvature, Analyse Nonlinéaire 9 (1992), 643-655.



**P.Amster and D. F. Rial**

Dpto. de Matemática Fac. de Cs. Exactas y Naturales, UBA Pab. I, Ciudad Universitaria (1428) Capital, Argentina

**M. C. Mariani**

Dpto. de Matemática Fac. de Cs. Exactas y Naturales, UBA Pab. I, Ciudad Universitaria (1428) Capital, Argentina

CONICET

**Address for correspondence:** Prof. M. C. Mariani, Dpto. de Matemática

Fac. de Cs. Exactas y Naturales, UBA Pab. I, Ciudad Universitaria (1428) Capital, Argentina

**E-mail: mcmarian@dm.uba.ar**