

THE OPTIMAL SIZE OF THE EXCHANGE HOLE AND REDUCTION TO ONE-PARTICLE HAMILTONIANS

PAUL MANCAS, A. M. KLAUS MÜLLER †, AND HEINZ SIEDENTOP

ABSTRACT. We use the concept of the exchange hole introduced by Slater to bound the energy of atoms, molecules, and other systems interacting by Coulomb forces from below by one-particle Hamiltonians with an effective screening potential and an exchange hole around each electron. Interestingly enough the optimal size of the exchange hole is smaller than Slater proposed: the best lower bound is obtained when the exchange hole carries charge 1/2 instead of 1. To highlight the quality of our estimate we show that the Dirac exchange energy with a slightly different constant bounds the exchange-correlation energy from below, an estimate previously derived by Lieb [5] and later improved by Lieb and Oxford [6].

1. INTRODUCTION

Slater [12] simplified the Hartree-Fock equations by introducing a ball around each electron that carries a unit charge modeling the self-interaction and the exchange energy of a system of N electrons. This ball, called “exchange hole” for brevity, replaces the exchange term in the Hartree-Fock equations and thus leads to a substantial simplification. Because of this importance it has attracted considerable interest since the pioneering work of Slater. We need to restrict ourselves for brevity but would like to mention the recent works of Buijse and Baerends [2], Springborg et al. [13], Becke and Roussel [1], and the references therein.

The novelty of our result is that it does not only give an expression for the exchange-correlation energy of atoms and molecules that becomes exact in the limit of large (neutral) atoms (see, e.g., Siedentop and Weikard [8, 8, 9, 10, 11]). It also yields a rigorous lower bound on the exact energy.

The observation relevant for us is due to Hughes [3, 4]. He noted that the classical Coulomb interaction of N point particles at positions $\mathbf{x}_1, \dots, \mathbf{x}_N$ with unit charge can be estimated from below as follows:

$$(1) \quad \sum_{1 \leq n < m \leq N} \frac{1}{|\mathbf{x}_n - \mathbf{x}_m|} \geq \sum_{n=1}^N \int_{|\mathbf{y} - \mathbf{x}_n| > R_1(\mathbf{x}_n)} \frac{\sigma(\mathbf{y}) d\mathbf{y}}{|\mathbf{y} - \mathbf{x}_n|} - D(\sigma, \sigma) + \sum_{n=1}^N \frac{1}{2R_1(\mathbf{x}_n)}$$

where we use the following notation: given any charge density σ in three dimensional space \mathbb{R}^3 we denote its electrostatic self-energy by $D(\sigma, \sigma)$, i.e.,

$$(2) \quad D(\sigma, \sigma) := \frac{1}{2} \int_{\mathbb{R}^3} d\mathbf{x} \int_{\mathbb{R}^3} d\mathbf{y} \frac{\sigma(\mathbf{x})\sigma(\mathbf{y})}{|\mathbf{x} - \mathbf{y}|}.$$

Date: September 19, 2003.

Key words and phrases. Exchange hole, correlation energy, exchange energy, Lieb-Oxford inequality, Coulomb systems.

The authors thank Walter Farkas and Christian Hainzl for helpful discussions. Financial support by the European Union under its IHP network program, grant HPRN-CT-2002-00277, is gratefully acknowledged.

Here and in the following we will assume that the above integral exists even when σ is replaced by its absolute value $|\sigma|$. We will also assume that $\int_{\mathbb{R}^3} d\mathbf{y} |\sigma(\mathbf{y})| / |\mathbf{x} - \mathbf{y}|$ is finite for almost all \mathbf{x} .

Furthermore, we fix any non-negative density and define $R_\xi(\mathbf{x})$ to be the radius of the smallest ball with center at \mathbf{x} containing charge ξ , i.e., the smallest R fulfilling

$$(3) \quad \int_{|\mathbf{y}-\mathbf{x}|\leq R} \sigma(\mathbf{y}) d\mathbf{y} = \xi.$$

We assume, of course, $0 < \xi < \int \sigma$. For $R_{1/2}(\mathbf{x})$ we write simply $R(\mathbf{x})$.

Let us note the following:

(i) The assumption that the exchange hole contains a unit charge was strongly influenced by Slater's heuristic arguments.

(ii) Although the inequality is true for any charge density σ , good results will be only obtained, if it is picked to be a realistic density. In fact, using the Thomas-Fermi density allows to obtain lower bounds that are correct up to second order in $Z^{-1/3}$ (Scott correction) (Siedentop and Weikard [9, 10, 11] and Hughes [3, 4]). – It is also possible to determine σ self-consistently, i.e., setting $\sigma := |u_1(x)|^2 + \dots + |u_N(x)|^2$ where the u_n are the solutions of the corresponding one-particle problem.

(iii) The Inequality (1) implies strong simplifications for the treatment of interacting systems: it allows to replace two-particle operators by one-particle operators, since their ground states are known to be Slater determinants. It is therefore of utmost importance to obtain a bound of this type that is as tight as possible.

The purpose of this paper is exactly this, i.e., to improve the above bound. We will no longer fix the amount of charge ξ in the exchange hole. Instead we will take it as a variational parameter. It will turn out that the bound on the interaction energy does not only simplify for $\xi = 1/2$; it will also improve and be tighter compared to all other values of ξ including 1.

As an application we will show in Section 3 that the resulting bound implies an inequality of Lieb [5] later improved by Lieb and Oxford [6].

2. AN OPTIMAL LOWER BOUND ON THE COULOMB INTERACTION VIA AN EXCHANGE HOLE

Our main result is the following inequality:

Theorem 1.

- (1) *Let σ be a charge density with $\int \sigma(\mathbf{y}) d\mathbf{y} \geq 1/2$. Then, for any given number N of points $\mathbf{x}_1, \dots, \mathbf{x}_N$ in space we have*

$$(4) \quad \sum_{1 \leq n < m \leq N} \frac{1}{|\mathbf{x}_n - \mathbf{x}_m|} \geq \sum_{n=1}^N \int_{|\mathbf{y}-\mathbf{x}_n| > R_\xi(\mathbf{x}_n)} \frac{\sigma(\mathbf{y})}{|\mathbf{y} - \mathbf{x}_n|} d\mathbf{y} - D(\sigma, \sigma) + \sum_{n=1}^N \frac{\xi - 1/2}{R_\xi(\mathbf{x}_n)}.$$

- (2) *The right hand side of (4) has its maximum for $\xi = 1/2$.*

We would like to remark that we can replace $\sigma(\mathbf{x}) d\mathbf{x}$ by any signed measure $d\mu(\mathbf{x})$ with the above requirements on its interaction energy and its potential. Our main result would still be true. We leave this to the interested reader.

Our proof depends on a well known formula that goes back to Newton [7]: for any spherically symmetric charge density ρ one has

$$(5) \quad \int_{\mathbb{R}^3} \frac{\rho(\mathbf{y}) d\mathbf{y}}{|\mathbf{x} - \mathbf{y}|} = \int_0^\infty \frac{4\pi r^2 \rho(r) dr}{\max\{r, |\mathbf{x}|\}}$$

which is easily verified by integration in spherical coordinates.

Proof. 1. It is convenient to introduce a charge density of total charge one which is smeared out uniformly on a sphere of radius R centered at $\mathbf{y} \in \mathbb{R}^3$:

$$(6) \quad d\mu_{R,\mathbf{y}}(\mathbf{x}) := \frac{\delta(|\mathbf{x} - \mathbf{y}| - R)d\mathbf{x}}{4\pi R^2}.$$

We begin with the crucial observation, that the Coulomb kernel is positive as an operator, i.e.,

$$(7) \quad D(\mu, \mu) \geq 0$$

for any complex measure μ on the three dimensional space provided the corresponding integral is absolute convergent, i.e., $\int_{\mathbb{R}^3} d|\mu|(\mathbf{x}) \int_{\mathbb{R}^3} d|\mu|(\mathbf{y}) |\mathbf{x} - \mathbf{y}|^{-1}$ is finite. Now, we choose

$$d\mu(x) := \sigma - d\mu_{R_\xi(\mathbf{x}_1), \mathbf{x}_1} - \dots - d\mu_{R_\xi(\mathbf{x}_N), \mathbf{x}_N}.$$

Rearranging the terms in (7) gives

$$(8) \quad 2 \sum_{1 \leq n < m \leq N} D(\mu_{R_\xi(\mathbf{x}_n), \mathbf{x}_n}, \mu_{R_\xi(\mathbf{x}_m), \mathbf{x}_m}) \\ \geq -D(\sigma, \sigma) - \sum_{n=1}^N D(\mu_{R_\xi(\mathbf{x}_n), \mathbf{x}_n}, \mu_{R_\xi(\mathbf{x}_n), \mathbf{x}_n}) + 2 \sum_{n=1}^N D(\sigma, \mu_{R_\xi(\mathbf{x}_n), \mathbf{x}_n}).$$

We evaluate and estimate the occurring expressions. By explicit computation using Newton's formula (5) we find

$$(9) \quad 2D(\mu_{R_\xi(\mathbf{x}_n), \mathbf{x}_n}, \mu_{R_\xi(\mathbf{x}_m), \mathbf{x}_m}) \leq |\mathbf{x}_n - \mathbf{x}_m|^{-1}$$

which is also obvious from the physical point of view: it costs energy to contract the smeared out unit charge to a point. This means, that we can estimate the left hand side of (8) from above by $\sum_{1 \leq n < m \leq N} |\mathbf{x}_n - \mathbf{x}_m|^{-1}$. Computing, the self-interaction of a unit charge smeared out on a sphere of radius R yields

$$(10) \quad D(\mu_{R,\mathbf{x}}, \mu_{R,\mathbf{x}}) = \frac{1}{2R}.$$

Finally, the summands of last term of the right side of (8) are again computed explicitly:

$$(11) \quad D(\sigma, \mu_{R_\xi(\mathbf{x}), \mathbf{x}}) = \frac{1}{2} \int_{|\mathbf{y}-\mathbf{x}| > R_\xi(\mathbf{x})} \frac{\sigma(\mathbf{y})}{|\mathbf{y} - \mathbf{x}|} d\mathbf{y} + \frac{\xi}{2R_\xi(\mathbf{x})}.$$

Inserting the estimates gives the desired first inequality

$$(12) \quad \sum_{1 \leq n < m \leq N} \frac{1}{|\mathbf{x}_n - \mathbf{x}_m|} \geq \sum_{n=1}^N \int_{|\mathbf{y}-\mathbf{x}_n| > R_\xi(\mathbf{x}_n)} \frac{\sigma(\mathbf{y})}{|\mathbf{y} - \mathbf{x}_n|} d\mathbf{y} - D(\sigma, \sigma) + \sum_{n=1}^N \frac{\xi - 1/2}{R_\xi(\mathbf{x}_n)}.$$

2. To prove the second part of the theorem we compare the right hand side *rhs* of (4) at $\xi = 1/2$ with all other values of $\xi = 1/2 + \epsilon$: we have – assuming $\epsilon > 0$ –

$$\begin{aligned}
(13) \quad rhs(1/2 + \epsilon) - rhs(1/2) &= \sum_{n=1}^N \int_{|\mathbf{y}-\mathbf{x}_n| > R_{1/2+\epsilon}(\mathbf{x}_n)} \frac{\sigma(\mathbf{y})}{|\mathbf{y}-\mathbf{x}_n|} d\mathbf{y} - D(\sigma, \sigma) \\
&+ \sum_{n=1}^N \frac{\epsilon}{R_{1/2+\epsilon}(\mathbf{x}_n)} - \sum_{n=1}^N \int_{|\mathbf{y}-\mathbf{x}_n| > R_{1/2}(\mathbf{x}_n)} \frac{\sigma(\mathbf{y})}{|\mathbf{y}-\mathbf{x}_n|} d\mathbf{y} + D(\sigma, \sigma) \\
&= - \sum_{n=1}^N \int_{R_{1/2}(\mathbf{x}_n) < |\mathbf{y}-\mathbf{x}_n| < R_{1/2+\epsilon}(\mathbf{x}_n)} \frac{\sigma(\mathbf{y})}{|\mathbf{y}-\mathbf{x}_n|} d\mathbf{y} + \sum_{n=1}^N \frac{\epsilon}{R_{1/2+\epsilon}(\mathbf{x}_n)} \\
&\leq - \sum_{n=1}^N \int_{R_{1/2}(\mathbf{x}_n) < |\mathbf{y}-\mathbf{x}_n| < R_{1/2+\epsilon}(\mathbf{x}_n)} \frac{\sigma(\mathbf{y})}{R_{1/2+\epsilon}(\mathbf{x}_n)} d\mathbf{y} + \sum_{n=1}^N \frac{\epsilon}{R_{1/2+\epsilon}(\mathbf{x}_n)} = 0,
\end{aligned}$$

since the annulus of integration contains exactly charge ϵ by definition of $R_\xi(x)$.

The case of negative ϵ is analogous. \square

To emphasize the charge optimized bound we state it explicitly:

$$(14) \quad \boxed{\sum_{1 \leq n < m \leq N} \frac{1}{|\mathbf{x}_n - \mathbf{x}_m|} \geq \sum_{n=1}^N \int_{|\mathbf{y}-\mathbf{x}_n| > R(\mathbf{x}_n)} \frac{\sigma(\mathbf{y}) d\mathbf{y}}{|\mathbf{y}-\mathbf{x}_n|} - D(\sigma, \sigma)}$$

which holds for arbitrary charge density σ fulfilling merely the general requirements of the introduction.

3. THE CORRELATION BOUND OF LIEB AND OXFORD

We would like to show that the Inequality (14) implies an exchange-correlation bound in terms of the Dirac exchange term that goes back to Lieb and Lieb and Oxford.

The main technical tool will be the Hardy-Littlewood maximal function $(Mf)(\mathbf{x})$ of a function f . It is defined to be the biggest spherical average of f over balls of radius R , i.e.,

$$(15) \quad (Mf)(\mathbf{x}) := \sup_{R>0} \frac{\int_{|\mathbf{x}-\mathbf{y}| < R} |f(\mathbf{y})| d\mathbf{y}}{\frac{4\pi}{3} R^3}.$$

It is a classical fact that the L^p norm of the maximal function can be estimated in terms of the L^p norm of the function itself (Stein [15]). We will need this fact in the case that $p = 4/3$. This estimate reads

$$(16) \quad \int_{\mathbb{R}^3} (Mf)(\mathbf{x})^{4/3} d\mathbf{x} \leq A_{4/3}^{4/3} \int_{\mathbb{R}^3} |f(\mathbf{x})|^{4/3} d\mathbf{x}.$$

(Note that best constant $A_{4/3}$ in this inequality is not exceeding 17.57 (Stein and Strömberg [14]). We also need an inequality that appears in the spherical symmetric case in Lieb [5]:

Lemma 1. *If f is an integrable function, then*

$$(17) \quad \int_{\mathbb{R}^3} |f(\mathbf{x})|/|\mathbf{x}| d\mathbf{x} \leq \left(\frac{9}{2} \pi (Mf)(\mathbf{0}) \right)^{1/3} \left(\int_{\mathbb{R}^3} |f(\mathbf{x})| d\mathbf{x} \right)^{2/3}.$$

Proof. Obviously, it is enough to prove the result for non-negative spherically symmetric functions. Integrating by parts we get

$$\begin{aligned}
(18) \quad \int_{\mathbb{R}^3} f(\mathbf{x})/|\mathbf{x}|d\mathbf{x} &= \frac{1}{4\pi} \int_{\mathbb{R}^3} d\mathbf{x}|\mathbf{x}|^{-4} \int_{|\mathbf{y}|<|\mathbf{x}|} d\mathbf{y}f(\mathbf{y}) \\
&= \frac{1}{3} \int_{|\mathbf{x}|<R} d\mathbf{x}|\mathbf{x}|^{-1} \frac{\int_{|\mathbf{y}|<|\mathbf{x}|} d\mathbf{y}f(\mathbf{y})}{\frac{4\pi}{3}|\mathbf{x}|^3} + \frac{1}{4\pi} \int_{|\mathbf{x}|>R} d\mathbf{x}|\mathbf{x}|^{-4} \int_{|\mathbf{y}|<|\mathbf{x}|} d\mathbf{y}f(\mathbf{y}) \\
&\leq \frac{2\pi}{3}R^2(Mf)(\mathbf{0}) + \int_{\mathbb{R}^3} d\mathbf{y}f(\mathbf{y})/R.
\end{aligned}$$

Picking $R = \sqrt[3]{3 \int_{\mathbb{R}^3} d\mathbf{y}f(\mathbf{y})/(4\pi(Mf)(\mathbf{0}))}$ yields the desired inequality. \square

Lemma 2. *Let σ be as required in the introduction. Then*

$$(19) \quad \int_{|\mathbf{x}-\mathbf{y}|\leq R(\mathbf{x})} d\mathbf{y} \frac{\sigma(\mathbf{y})}{|\mathbf{x}-\mathbf{y}|} \leq \frac{(9\pi)^{1/3}}{2} (M\sigma)(\mathbf{x})^{1/3}.$$

Proof. We set $\sigma_{\mathbf{x}}(\mathbf{y}) := \sigma(\mathbf{x} + \mathbf{y})\chi_{R(\mathbf{x})}(\mathbf{y})$ where $\chi_{R(\mathbf{x})}(\mathbf{y})$ is one, if $|\mathbf{y}| < R(\mathbf{x})$ and vanishes otherwise. We apply Lemma 1 with $f = \sigma_{\mathbf{x}}$ and get

$$\begin{aligned}
(20) \quad \int_{|\mathbf{y}-\mathbf{x}|<R(\mathbf{x})} d\mathbf{y} \frac{\sigma(\mathbf{y})}{|\mathbf{x}-\mathbf{y}|} &= \int_{\mathbb{R}^3} d\mathbf{y} \frac{\sigma_{\mathbf{x}}(\mathbf{y})}{|\mathbf{y}|} \\
&\leq \left(\frac{9}{2}\pi(M\sigma_{\mathbf{x}})(\mathbf{0}) \right)^{1/3} \left(\int_{\mathbb{R}^3} |\sigma_{\mathbf{x}}(\mathbf{y})|d\mathbf{y} \right)^{2/3} = \left(\frac{9}{2}\pi(M\sigma)(\mathbf{x}) \right)^{1/3} \left(\frac{1}{2} \right)^{2/3}
\end{aligned}$$

where the last inequality holds because of the definition of $R(\mathbf{x})$. The last term of the above chain of inequalities is easily seen to give the claim. \square

Assume ψ to be a normalized wave function of N particles with spin q each, i.e., ψ depends on the variables $\vec{x} := (x_1, \dots, x_N)$ where each $x_n = (\mathbf{x}_n, \tau_n)$ is a space-spin variable, i.e.,

$$\int |\psi(\vec{x})|^2 d\vec{x} := \int_{\mathbb{R}^3} d\mathbf{x}_1 \sum_{\tau_1=1}^q \dots \int_{\mathbb{R}^3} d\mathbf{x}_N \sum_{\tau_N=1}^q |\psi(\vec{x})|^2 = 1.$$

We do not assume that ψ fulfills any symmetry requirements but we do assume that

$$I_{\psi} := \sum_{1 \leq n < m \leq N} \int d\vec{x} \frac{|\psi(\vec{x})|^2}{|\mathbf{x}_n - \mathbf{x}_m|} < \infty$$

and that the one-particle density ρ_{ψ} of the state ψ is integrable when raised to the power $4/3$, i.e., $\int \rho_{\psi}^{4/3} < \infty$. Note that all these requirements are naturally fulfilled for quantum states with finite kinetic energy. Then

$$(21) \quad I_{\psi} \geq 2D(\rho_{\psi}, \sigma) - D(\sigma, \sigma) - L \left(\int_{\mathbb{R}^3} \rho_{\psi}(\mathbf{x})^{4/3} d\mathbf{x} \right)^{3/4} \left(\int_{\mathbb{R}^3} \sigma(\mathbf{x})^{4/3} d\mathbf{x} \right)^{1/4}$$

with $L = (9\pi)^{1/3} A_{4/3}^{4/9}/2$. (Using the above upper bound of Stein and Strömberg on $A_{4/3}$ shows that $L \leq 3.96$.) Picking $\sigma = \rho_{\psi}$ yields the inequality of Lieb [5] and Lieb and Oxford [6]

$$(22) \quad I_{\psi} \geq D(\rho_{\psi}, \rho_{\psi}) - L \int_{\mathbb{R}^3} \rho_{\psi}(\mathbf{x})^{4/3} d\mathbf{x}.$$

Note that Lieb [5] estimated L by 8.52 using also the maximal inequality and Lieb and Oxford [6] estimated L by 1.68 using a somewhat more involved but elementary technique. That our result implies this inequality with a relatively good constant

highlights the quality of our exchange-correlation estimate using the idea of the exchange hole.

We start the proof of Inequality (21) by multiplying both sides of Inequality (14) by $|\psi(\vec{x})|^2$ and integrate and sum over all space-spin variables:

$$(23) \quad I_\psi \geq 2D(\rho_\psi, \sigma) - \int_{\mathbb{R}^3} d\mathbf{x} \rho_\psi(\mathbf{x}) \int_{|\mathbf{x}-\mathbf{y}| < R(\mathbf{x})} \frac{\sigma(\mathbf{y}) d\mathbf{y}}{|\mathbf{x}-\mathbf{y}|} - D(\sigma, \sigma) \\ \geq 2D(\rho_\psi, \sigma) - D(\sigma, \sigma) - \frac{(9\pi)^{1/3}}{2} \int_{\mathbb{R}^3} \rho_\psi(\mathbf{x}) (M\sigma)(\mathbf{x})^{1/3} d\mathbf{x}$$

where we have used (19) of Lemma 2. Applying the Hölder inequality followed by the maximal inequality (16) gives

$$(24) \quad I_\psi \geq 2D(\rho_\psi, \sigma) - D(\sigma, \sigma) - \frac{(9\pi)^{1/3}}{2} \left(\int_{\mathbb{R}^3} \rho_\psi^{4/3} \right)^{3/4} \left(\int_{\mathbb{R}^3} (M\sigma)^{4/3} \right)^{1/4} \\ \geq 2D(\rho_\psi, \sigma) - D(\sigma, \sigma) - \frac{(9\pi)^{1/3}}{2} A_{4/3}^{4/9} \left(\int_{\mathbb{R}^3} \rho_\psi^{4/3} \right)^{3/4} \left(\int_{\mathbb{R}^3} \sigma^{4/3} \right)^{1/4}$$

which proves Inequality (21).

4. APPROXIMATE DENSITY FUNCTIONAL

The Inequality (14) reduces the interacting N -particle problem to an approximate non-interacting one whose energy is a strict lower bound on the exact energy. Picking in addition the occurring arbitrary density σ appropriately, allows us to construct a Kohn-Sham type density functional for the ground state energy which is not only an approximation but gives a rigorous lower bound: we denote the Kohn-Sham orbitals of the N electrons by ϕ_1, \dots, ϕ_N . Next we pick

$$\sigma(\mathbf{x}) := \rho(\mathbf{x}) := |\phi_1(\mathbf{x})|^2 + \dots + |\phi_N(\mathbf{x})|^2.$$

The approximate Kohn-Sham functional becomes in this case

$$\mathcal{E}_{KS}[\phi_1, \dots, \phi_N] := \sum_{\nu=1}^N \int |\nabla \phi_\nu|^2 + D(\rho, \rho) - \int d\mathbf{x} \int d\mathbf{y}_{|\mathbf{y}-\mathbf{x}| < R(\mathbf{x})} \frac{\rho(\mathbf{x})\rho(\mathbf{y})}{|\mathbf{x}-\mathbf{y}|} \\ + \int V(\mathbf{x})\rho(\mathbf{x})d\mathbf{x}$$

where V is the external electric potential. The corresponding Kohn-Sham equations are

$$-\Delta \phi_\nu + \left(V(\mathbf{x}) + \int_{|\mathbf{x}-\mathbf{y}| > R(\mathbf{x})} \frac{\rho(\mathbf{y})d\mathbf{y}}{|\mathbf{x}-\mathbf{y}|} \right) \phi_\nu = \epsilon_\nu \phi_\nu.$$

REFERENCES

- [1] A. D. Becke and M. R. Roussel. Exchange holes in inhomogeneous systems – a coordinate-space model. *Phys. Rev. A*, 39(8):3761–3767, April 1989.
- [2] M. A. Buijse and Baerends E. J. An approximate exchange-correlation hole density as a functional of the natural orbitals. *Molecular Physics*, 100(4):401–421, February 2002.
- [3] Webster Hughes. *An Atomic Energy Lower Bound that Gives Scott's Correction*. PhD thesis, Princeton, Department of Mathematics, 1986.
- [4] Webster Hughes. An atomic lower bound that agrees with Scott's correction. *Adv. in Math.*, 79:213–270, 1990.
- [5] Elliott H. Lieb. A lower bound for Coulomb energies. *Phys. Lett.*, 70A:444–446, 1979.
- [6] Elliott H. Lieb and Stephen Oxford. Improved lower bound on the indirect Coulomb energy. *Intern. J. Quantum Chem.*, 19:427–439, 1981.

- [7] Isaac Newton. *Philosophiae naturalis principia mathematica. Vol. I.* Harvard University Press, Cambridge, Mass., 1972. Reprinting of the third edition (1726) with variant readings, Assembled and edited by Alexandre Koyré and I. Bernard Cohen with the assistance of Anne Whitman.
- [8] Heinz Siedentop and Rudi Weikard. Upper bound on the ground state energy of atoms that proves Scott's conjecture. *Phys. Lett. A*, 120:341–342, 1987.
- [9] Heinz Siedentop and Rudi Weikard. On the leading energy correction of the statistical atom: Lower bound. *Europhysics Letters*, 6:189–192, 1988.
- [10] Heinz Siedentop and Rudi Weikard. On the leading correction of the Thomas-Fermi model: Lower bound – with an appendix by A. M. K. Müller. *Invent. Math.*, 97:159–193, 1989.
- [11] Heinz Siedentop and Rudi Weikard. A new phase space localization technique with application to the sum of negative eigenvalues of Schrödinger operators. *Annales Scientifiques de l'École Normale Supérieure*, 24(2):215–225, 1991.
- [12] John C. Slater. A simplification of the Hartree-Fock method. *Physical Review*, 81(3):385–390, February 1951.
- [13] M. Springborg, J. P. Perdew, and K. Schmidt. Phase space, density matrices, energy densities, and exchange holes. *Zeitschrift für Physikalische Chemie*, 215:1243–1264, 2000.
- [14] E. M. Stein and J.-O. Strömberg. Behavior of maximal functions in \mathbf{R}^n for large n . *Ark. Mat.*, 21(2):259–269, 1983.
- [15] Elias M. Stein. *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, volume 43 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1993. With the assistance of Timothy S. Murphy, Monographs in Harmonic Analysis, III.

MATHEMATISCHES INSTITUT, THERESIENSTR. 39, 80333 MÜNCHEN, GERMANY

E-mail address: paul_mancas@yahoo.de

INSTITUT FÜR MATHEMATISCHE PHYSIK, TECHNISCHE UNIVERSITÄT BRAUNSCHWEIG, 38106
BRAUNSCHWEIG, GERMANY

MATHEMATISCHES INSTITUT, THERESIENSTR. 39, 80333 MÜNCHEN, GERMANY

E-mail address: h.s@lmu.de