

Periodic Nonlinear Schrödinger Equation with Application to Photonic Crystals

ALEXANDER PANKOV
Mathematics Department
College of William and Mary
Williamsburg, VA 23187-8795
e-mail: `pankov@member.ams.org`

Abstract

We present basic results, known and new, on nontrivial solutions of periodic stationary nonlinear Schrödinger equations. We also sketch an application to nonlinear optics and discuss some open problems.

0 Introduction

We consider the nonlinear Schrödinger equation (NLS)

$$-\Delta u + V(x)u = \pm f(x, u), \quad x \in \mathbb{R}^N, \quad (1)$$

with superlinear subcritical nonlinearity f , $f(x, 0) = 0$. Our basic assumption is periodicity of $V(x)$ and $f(x, u)$ in x with respect to some lattice $\Lambda \subset \mathbb{R}^N$. For the sake of simplicity in the following we assume that $\Lambda = \mathbb{Z}^N$ is the standard integer lattice in \mathbb{R}^N , i.e. the functions $V(x)$ and $f(x, u)$ are 1-periodic in each variable x_i , $x = (x_1, x_2, \dots, x_N)$. We are interested in nontrivial solutions of equation (1) such that $u(\infty) = 0$.

Problems of such kind appear in many applications. For instance, in condensed matter physics, the states of atomic gas are described by wave functions $\psi(t, x)$ that satisfy the Gross-Pitaevski equation (see, e.g., [14])

$$i\psi_t = -\Delta\psi + V_0(x)\psi - \alpha|\psi|^2\psi$$

(the so-called mean field approximation). Condensed states confined in a finite region (Bose-Einstein condensates) correspond to the wave functions $\psi = u(x) \cdot \exp(-i\omega t)$ where the amplitude $u(x)$ vanishes at infinity, i.e. standing waves. Here $V_0(x)$ is the external

electric potential which can be periodic. Making use the standing wave Ansatz, we arrive at the equation of the form (1). Another field of applications, gap solitons in photonic crystals, will be discussed in Section 7 below.

There is a number of papers dealing with periodic NLS and its solutions vanishing at infinity. First, the case when the operator $-\Delta + V$ is positive definite was considered. In [24] the author has proved an existence theorem using the Nehari variational principle (see, e.g., [38]) and concentrated compactness results [20]. (Even more general asymptotically periodic case was treated in that paper). Later P. H. Rabinowitz [30] has obtained the existence of nontrivial solutions under less restrictive assumptions on the nonlinearity f . Moreover, in [13] V. Coti Zelati and P. H. Rabinowitz have proved the existence of infinitely many multi-bump solutions, imposing some condition of general position.

In general, it is known (see, e.g., [32]) that the spectrum of the self-adjoint operator $-\Delta + V$ in $L^2(\mathbb{R}^N)$ is purely continuous and may contain gaps, i.e. open intervals free of spectrum. So, it is possible that 0 is in a finite spectral gap and the operator $-\Delta + V$ is not positive definite. In this case first existence results (under very strong assumptions on the nonlinearity) were found in [4, 15]. Later C. Troestler and M. Willem [37] and W. Kryszewski and A. Szulkin [17] have proved the existence of nontrivial solutions under much more natural conditions. Their proofs are based on a generalized linking theorems applied to the corresponding action functional. The original proofs of such linking results required new degree theories in order to overcome the lack of compactness and find at least one Palais-Smale sequence. Moreover, in [17] it is shown that there exists infinitely many geometrically distinct solutions, provided the nonlinearity is odd. It is not known whether these solutions are multibumps. In [7], T. Bartsch and Y. Ding considered the case when 0 is a boundary point of the spectrum and, in addition, gave a simpler proof the generalized linking theorem. In [25], the author and K. Pflüger suggested another approach to the periodic NLS based on periodic approximations and standard critical point theorems. This approach goes back to P. H. Rabinowitz [30] who considered the positive definite case. Later the same idea was applied to other problems [26, 27, 28]. See also [1, 4, 5, 15, 11, 19, 39] for some other results on NLS.

Now we list the basic assumptions. Let

$$F(x, u) = \int_0^u f(x, s) ds.$$

- (i) $V \in \infty(\mathbb{R}^N)$ is a 1-periodic function in each x_i , $i = 1, \dots, N$.
- (ii) $f : \mathbb{R}^N \times \mathbb{R} \rightarrow \mathbb{R}$ is a Carathéodory function¹ that is 1-periodic in each x_i , $i = 1, \dots, N$.
- (iii) $|f(x, u)| \leq C(1 + |u|^{p-1})$ where $2 < p < 2^*$, $2^* := 2N/(N-2)$ if $N \geq 3$ and $2^* := +\infty$ if $N = 1, 2$.
- (iv)

$$\operatorname{ess\,sup}_{x \in \mathbb{R}^N} |f(x, u)| = o(|u|), \quad u \rightarrow 0.$$

¹This means that $f(x, u)$ is measurable in $x \in \mathbb{R}^N$ for all $u \in \mathbb{R}$ and continuous in $u \in \mathbb{R}$ for a.e. $x \in \mathbb{R}^N$.

(v) *There exists $q > 2$ such that*

$$0 < qF(x, u) \leq uf(x, u), \quad u \neq 0.$$

Assumption (i) implies that the operator $-\Delta + V$ with the domain $C_0^\infty(\mathbb{R}^N)$ is essentially self-adjoint in $L^2(\mathbb{R}^N)$. Denote by L its self-adjoint extension and $\sigma(L)$ the spectrum of L . The last assumption is the following

(vi) *0 is not in $\sigma(L)$. In addition, in the case of ‘-’ in equation (1) we assume that $\inf \sigma(L) < 0$.*

This assumption implies that there is a maximal open interval $(-\alpha_-, \alpha_+)$, with $\alpha_\pm > 0$, free of the spectrum (gap in the spectrum). In the case of ‘+’ sign it is possible that $\alpha_- = \infty$. However, in the case of ‘-’ sign both α_- and α_+ are finite, i.e. 0 is in a finite gap. We set $\alpha := \min[\alpha_+, \alpha_-]$. Note that α is the distance from 0 to $\sigma(L)$.

The power nonlinearity $f(x, u) = h(x)|u|^{p-2}u$, with positive 1-periodic $h \in L^\infty(\mathbb{R}^N)$ and $2 < p < 2^*$, satisfies all the assumptions. The cubic nonlinearity ($p = 4$) is admissible if $N = 1, 2$ or 3 . The last nonlinearity is important for many applications (see, e.g., Section 7). Note that in the case $N = 1$ assumption (iii) can be replaced by the following one. For any $h > 0$ there exists a constant $C(h) > 0$ such that

$$\operatorname{ess\,sup}_{x \in \mathbb{R}^N} |f(x, u)| \leq C(h) \quad \text{if } |u| \leq h.$$

Assumptions (iii) and (v) mean that we consider subcritical superlinear nonlinearities. For some results for the critical periodic NLS we refer to [12]. Another important problem concerns the existence of solutions to (1) vanishing at infinity in the case of asymptotically linear nonlinearity (see a remark in Section 7). Some results in this direction can be found in [19], but the problem is still not well understood.

As it follows from (v), the nonlinearity $f(x, u)$ does not change sign when x changes. In the case of periodic NLS with sign changing nonlinearity the existence of vanishing at infinity solutions is a completely open problem that may have interesting consequences in the theory of gap solitons (see discussion in Section 7).

Now we summarize the principal result.

Theorem 1 *Under assumptions (i)–(vi) equation (1) has a nontrivial weak solution $u \in H^1(\mathbb{R}^N)$. This solution is continuous and decays exponentially at infinity.*

In subsequent sections we present the proof and discuss some related results. We give also an application to the problem of existence of gap solitons in photonic crystals.

Throughout the paper we deal basically with the case of ‘+’ sign in (1) and only point out minor changes needed to cover the case of ‘-’ sign.

Acknowledgments. This article was prepared during author’s staying at Texas A&M University and College of William & Mary as a visiting professor. The author also thanks T. Bartsch, P. Kuchment, P. H. Rabinowitz, A. Szulkin and I. Spitkovsky for many interesting discussions. The work is partially supported by NATO, grant 970179.

1 Abstract critical point theorems

Let H be a real Hilbert space with the inner product (\cdot, \cdot) and the norm $\|\cdot\|$, and J a functional on H of the class $C^1(H)$. Denote by $J'(u)$ the Gateaux derivative (gradient) of J at the point $u \in H$. Recall that $u \in H$ is a *critical point* of J if $J'(u) = 0$ and $c = J(u)$ is the corresponding *critical value*. We shall use the following Palais-Smale condition:

(PS) Any sequence $u_n \in H$ such that $J(u_n)$ is bounded and $J'(u_n) \rightarrow 0$, i.e. Palais-Smale sequence, has a convergent subsequence.

Now we present the Linking Theorem (see [29, 38] for the proof). Let $H = Y \oplus Z$, $\rho > r > 0$ and let $z \in Z$ be such that $\|z\| = r$. Define

$$M := \{u = y + \lambda z : \|u\| \leq \rho, \lambda \geq 0, y \in Y\},$$

$$M_0 := \partial M = \{u = y + \lambda z : y \in Y, \|u\| = \rho \text{ and } \lambda \geq 0, \text{ or } \|u\| \leq \rho \text{ and } \lambda = 0\},$$

$$N := \{u \in Z : \|u\| = r\}.$$

We say that the functional J on H possesses the *Linking Geometry* if

$$b := \inf_{u \in N} J(u) > a := \sup_{u \in M_0} J(u).$$

Theorem 2 Suppose J satisfies (PS) and possesses the Linking Geometry with $\dim Y < \infty$. Let

$$c := \inf_{\gamma \in \Gamma} \sup_{u \in M} J(\gamma(u))$$

where

$$\Gamma := \{\gamma \in C(M, H) : \gamma = \text{id on } M_0\}.$$

Then c is a critical value of J and $b \leq c \leq \sup_{u \in M} J(u)$.

Note that in the case $Y = \{0\}$ Theorem 2 reduces to the Mountain Pass Theorem (see, e.g., [29, 38]).

Sometimes one can drop the assumption $\dim Y < \infty$ in the Linking Theorem. The following result was obtained in [9].

Theorem 3 Assume that the functional J possesses the Linking Geometry and satisfies the following assumptions

(j) $J(u) = \frac{1}{2}(Au, u) + b(u)$ where $u = u_1 + u_2 \in Y \oplus Z$, $Au = A_1u_1 + A_2u_2$ $A_1 : Y \rightarrow Y$ and $A_2 : Z \rightarrow Z$ are linear bounded self-adjoint operators;

(jj) b is weakly continuous and uniformly differentiable on bounded sets;

(jjj) Any sequence $u_n \in H$ such that $J(u_n)$ is bounded from above and $J'(u_n) \rightarrow 0$ is bounded.

Then c defined in Theorem 2 is a critical value of J such that $b \leq c \leq \sup_{u \in M} J(u)$.

Note that in the case of Theorem 3 the functional J satisfies the Palais-Smale condition.

2 Variational setting and periodic problem

Throughout this section we will suppose assumptions (i)–(vi) to be valid.

Let Q_k be the cube in \mathbb{R}^N , with the edge length k , centered at the origin. Denote by L_k a unique self-adjoint operator in $L^2(Q_k)$ generated by $-\Delta + V$ with periodic boundary conditions. We can (and will consider) this space as the space of all k -periodic functions from $L^2_{\text{loc}}(\mathbb{R}^N)$. Note that $\sigma(L_k)$ is discrete and is a subset of $\sigma(L)$ (see, e.g., [32]). Let $E_k = H^1_{\text{per}}(Q_k)$ be the subspace of $H^1_{\text{loc}}(\mathbb{R}^N)$ that consists of k -periodic functions. This is just the form domain of L_k . Endowed with the standard $H^1(Q_k)$ norm, E_k is a Hilbert space. We also set $E = H^1(\mathbb{R}^N)$. This space coincides with the form domain of L .

Let E_k^+ (respectively, E_k^-) be the intersection of E_k with the positive (respectively, negative) spectral subspace of L_k . Similarly we introduce the subspaces E^+ and E^- in E . It is easily seen that $E_k^+ = (E_k^-)^\perp$ and $E^+ = (E^-)^\perp$. Thus, $E_k = E_k^- \oplus E_k^+$ and $E = E^- \oplus E^+$. These decompositions define the orthogonal projections P_k^\pm onto E_k^\pm and P^\pm onto E^\pm . For the sake of convenience, we introduce new inner products $(\cdot, \cdot)_k$ and (\cdot, \cdot) in E_k and E , with corresponding norms $\|\cdot\|_k$ and $\|\cdot\|$, so that

$$\int_{Q_k} (|\nabla u|^2 + V(x)|u|^2) dx = -\|u\|_k^2 \quad \text{for } u \in E_k^-,$$

$$\int_{Q_k} (|\nabla u|^2 + V(x)|u|^2) dx = \|u\|_k^2 \quad \text{for } u \in E_k^+$$

and

$$\int_{\mathbb{R}^N} (|\nabla u|^2 + V(x)|u|^2) dx = -\|u\|^2 \quad \text{for } u \in E^-,$$

$$\int_{\mathbb{R}^N} (|\nabla u|^2 + V(x)|u|^2) dx = \|u\|^2 \quad \text{for } u \in E^+.$$

These norms are equivalent to the standard H^1 -norms independently of k , i.e. there exists a constant $C > 0$, independent of k , such that

$$\alpha^{1/2}\|u\|_{H^1(Q_k)} \leq \|u\|_k \leq C\|u\|_{H^1(Q_k)}$$

and

$$\alpha^{1/2}\|u\|_{H^1(\mathbb{R}^N)} \leq \|u\| \leq C\|u\|_{H^1(\mathbb{R}^N)}.$$

On the subspaces E_k^\pm and E^\pm the constant α can be replaced by α_\pm . Moreover, the constant C can be chosen independent of V when V runs a bounded subset of $L^\infty(\mathbb{R}^N)$.

Consider the action functionals

$$\begin{aligned} J_k(u) &= \frac{1}{2}(\|P_k^+ u\|_k^2 - \|P_k^- u\|_k^2) \mp \int_{Q_k} F(x, u) dx = \\ &= \frac{1}{2} \int_{Q_k} (|\nabla u|^2 + V(x)|u|^2 \mp F(x, u)) dx \end{aligned}$$

and

$$\begin{aligned} J(u) &= \frac{1}{2}(\|P^+u\|^2 - \|P^-u\|^2) \mp \int_{\mathbb{R}^N} F(x, u) dx = \\ &= \frac{1}{2} \int_{\mathbb{R}^N} (|\nabla u|^2 + V(x)|u|^2 \mp F(x, u)) dx \end{aligned}$$

on E_k and E , respectively. From the standard Sobolev embedding theorem (see, e.g., [2, 21]) it follows that these functionals are well defined. One can also check that they are of the class C^1 and the derivatives are given by the formulas

$$\begin{aligned} \langle J'_k(u), v \rangle &= (P_k^+u, v)_k - (P_k^-u, v)_k \mp \int_{Q_k} f(x, u)v dx = \\ &= \int_{Q_k} (\nabla u \cdot \nabla v + V(x)uv \mp f(x, u)v) dx, \quad \forall v \in E_k \end{aligned}$$

and

$$\begin{aligned} \langle J'(u), v \rangle &= (P^+u, v) - (P^-u, v) \mp \int_{\mathbb{R}^N} f(x, u)v dx = \\ &= \int_{\mathbb{R}^N} (\nabla u \cdot \nabla v + V(x)uv \mp f(x, u)v) dx \quad \forall v \in E. \end{aligned}$$

Therefore, critical points of these functionals are weak solutions of (1), k -periodic in the case of J_k and vanishing, in a sense, at infinity in the case of J .

Recall that the gradient $\nabla J_k(u) \in E_k$ (resp., $\nabla J(u) \in E$) is defined by $(\nabla J_k(u), v)_k = \langle J'_k(u), v \rangle$ (resp., $(\nabla J(u), v)_k = \langle J'(u), v \rangle$) for all $v \in E_k$ (resp., $v \in E$).

Lemma 1 *There exists a constant $C > 0$ independent on k such that for any nontrivial critical points $u_k \in E_k$ of J_k and $u \in E$ of J , with critical values c_k and c respectively, we have*

$$\|u_k\|_k \leq C(|c_k|^{1/2} + |c_k|^{1/p'}), \quad \|u\| \leq C(|c|^{1/2} + |c|^{1/p'})$$

where $p' := p/(p-1)$ is the conjugate exponent.

Proof. From $c_k = J_k(u_k)$ and $J'_k(u_k) = 0$ we obtain

$$\begin{aligned} |c_k| &= |J_k(u_k) - \frac{1}{2} \langle J'_k(u_k), u_k \rangle| \geq \\ &\geq \left(\frac{1}{2} - \frac{1}{q} \right) \int_{Q_k} f(x, u_k)u_k dx. \end{aligned} \tag{2}$$

From assumptions (iii)–(v) we have the inequalities

$$|f(x, u)|^2 \leq Cuf(x, u) \quad \text{if } |u| \leq 1, \tag{3}$$

$$|f(x, u)|^{p'} \leq C|u|^{(p-1)(p'-1)} f(x, u) = Cuf(x, u) \quad \text{if } |u| \geq 1. \quad (4)$$

Let

$$B_k = \{x \in Q_k \mid |u_k(x)| \leq 1\}$$

Then from (2)–(4) we obtain

$$|c_k| \geq C' \left(\int_{B_k} |f(x, u_k(x))|^2 dx + \int_{Q_k \setminus B_k} |f(x, u_k(x))|^{p'} dx \right) \quad (5)$$

where $C' = (q-2)/(2qC)$. Hence

$$I_1 := \int_{B_k} |f(x, u_k(x))|^2 dx \leq |c_k|/C',$$

$$I_2 := \int_{Q_k \setminus B_k} |f(x, u_k(x))|^{p'} dx \leq |c_k|/C'.$$

Let $y_k = P_k^- u_k$ and $z_k = P_k^+ u_k$. From $\langle J'_k(u_k), y_k \rangle = 0$ and Hölder's inequality we see that

$$\|y_k\|_k^2 = - \int_{Q_k} f(x, u_k) y_k dx \leq I_1^{1/2} \|y_k\|_2 + I_2^{1/2} \|y_k\|_{p'}$$

where $\|\cdot\|_r$ stands for the standard L^r -norm. Then the Sobolev embedding theorem implies

$$\|y_k\|_k^2 \leq C'' \left(I_1^{1/2} + I_2^{1/p'} \right) \|y_k\|_k.$$

Clearly, the same argument works for z_k and we obtain

$$\|z_k\|_k^2 \leq C'' \left(I_1^{1/2} + I_2^{1/p'} \right) \|z_k\|_k.$$

These two inequalities imply the first estimate of the lemma. Similarly, we obtain the second one. \square

Remark 1 Similarly, we obtain the following more precise estimates

$$\alpha^{1/2} \|u_k\|_{H^1(Q_k)} \leq C(|c_k|^{1/2} + |c_k|^{1/p'}), \quad \alpha^{1/2} \|u\|_{H^1(\mathbb{R}^N)} \leq C(|c|^{1/2} + |c|^{1/p'}).$$

Lemma 2 *There exist constants $\varepsilon_1 > 0$ and $\varepsilon_2 > 0$ independent on k such that for any nontrivial critical points $u_k \in E_k$ of J_k and $u \in E$ of J we have $\|u_k\|_k \geq \varepsilon_1$, $\|u\| \geq \varepsilon_1$, $\mp J_k(u_k) \geq \varepsilon_2$ and $\pm J(u) \geq \varepsilon_2$.*

Proof. Consider ‘+’ sign case. Assumptions (iii) and (iv) implies immediately that for any $\varepsilon > 0$ there exists $C_\varepsilon > 0$ such that

$$|f(x, t)| \leq \varepsilon|t| + C_\varepsilon|t|^{p-1}.$$

Since $J'_k(u_k) = 0$, then $\langle J'_k(u_k), v_k \rangle = 0$ for any $v_k \in E_k$. Take $v_k = u_k^+ = P_k^+ u_k$. Then we have

$$\begin{aligned} \|u_k^+\|_k^2 &= \int_{Q_k} f(x, u_k) u_k^+ dx \leq \\ &\leq C(\varepsilon \|u_k\|_{L^2(Q_k)} \|u_k^+\|_{L^2(Q_k)} + C_\varepsilon \|u_k\|_{L^p(Q_k)}^{p-1} \|u_k^+\|_{L^p(Q_k)}) \\ &\leq C(\varepsilon \|u_k\|_k \|u_k^+\|_k + C_\varepsilon \|u_k\|_k^{p-1} \|u_k^+\|_k) \end{aligned}$$

where we have used the Sobolev embedding theorem. Similarly, taking $v_k = u_k^- = P_k^- u_k$ we obtain

$$\begin{aligned} \|u_k^-\|_k^2 &= \int_{Q_k} f(x, u_k) u_k^- dx \leq \\ &\leq C(\varepsilon \|u_k\|_k \|u_k^-\|_k + C_\varepsilon \|u_k\|_k^{p-1} \|u_k^-\|_k). \end{aligned}$$

These two inequalities imply immediately

$$\|u_k\|_k^2 = \|u_k^+\|_k^2 + \|u_k^-\|_k^2 \leq C(\varepsilon \|u_k\|_k^2 + C_\varepsilon \|u_k\|_k^p).$$

If ε is small enough, we obtain

$$\|u_k\|_k \geq \left(\frac{1 - \varepsilon C}{C_\varepsilon} \right)^{1/(p-2)} =: \varepsilon_1 > 0.$$

The same argument, with the same choice of constants, works for J . The estimates for critical values follow from Lemma 1.

The case of ‘-’ sign is similar. □

Theorem 4 *The functional J_k has a nontrivial critical point $u_k \in E_k$ with critical value c_k . Moreover, there exist a constant $C > 0$ independent of k such that $\|u_k\|_k \leq C$ and $0 < \pm c_k \leq C$.*

Proof. We start with the Palais-Smale condition. Refining the proof of Lemma 1, one can show that any Palais-Smale sequence for J_k is bounded in E_k . Now, using the standard argument [29, 38] based on the compactness of Sobolev embedding, we conclude that such a sequence is compact in E_k .

Consider ‘+’ sign case. We show that the functional J_k possesses the linking geometry with $Y = E_k^-$ and $Z = E_k^+$.

For $z \in Z$ we have

$$J_k(z) = \frac{1}{2} \|z\|_k^2 - \int_{Q_k} F(x, z) dx.$$

Assumptions (iii) and (iv) imply that for any $\varepsilon > 0$ there exists a constant $C_\varepsilon > 0$ such that $0 \leq F(x, u) \leq \varepsilon |u|^2 + C_\varepsilon |u|^p$. By the Sobolev embedding theorem we obtain

$$\int_{Q_k} F(x, z) dx \leq C(\varepsilon \|z\|_k^2 + C_\varepsilon \|z\|_k^p).$$

Choosing an appropriate ε we see that, for some $\delta > 0$, $J_k(z) \geq \delta$ if $\|z\|_k = r$ is small enough. Thus, we have chosen the sphere $N = \{u \in Z : \|u\|_k = r\}$.

Let us now find M . Fix any $z^0 \in Z = E_k^+$, $\|z^0\|_k = 1$. For $y + tz^0 \in M$, $y \in Y = E_k^-$ we have

$$J_k(y + tz^0) = \frac{1}{2}t^2 - \frac{1}{2}\|y\|_k^2 - \int_{Q_k} F(x, y + tz^0) dx.$$

Assumptions (iv) and (v) imply that for any $\varepsilon > 0$ there exists a constant $C_\varepsilon > 0$ such that $F(x, u) \geq -\varepsilon|u|^2 + C_\varepsilon|u|^q$. Hence,

$$\int_{Q_k} F(x, y + tz^0) dx \geq -\varepsilon\|y\|_{L^2(Q_k)}^2 - \varepsilon\|z^0\|_{L^2(Q_k)}^2 + C_\varepsilon\|y + tz^0\|_{L^q(Q_k)}^q.$$

Applying the Sobolev embedding theorem, we obtain

$$J_k(y + tz^0) \leq -\frac{1}{2}\|y\|_k^2 + \varepsilon C\|y\|_k^2 + \frac{1}{2}t^2 + \varepsilon C t^2 - C_\varepsilon\|y + tz^0\|_{L^q(Q_k)}^q. \quad (6)$$

Consider the space $X = Y \oplus \mathbb{R}z^0$ equipped with the $L^q(Q_k)$ -norm. Then $y + tz^0 \mapsto tz^0$ is a bounded projector in X . Since its norm is not less than 1, we see that

$$\|y + tz^0\|_{L^q(Q_k)} \geq \|tz^0\|_{L^q(Q_k)}.$$

Choosing ε so small that $\varepsilon C = 1/4$, we obtain from (6)

$$J_k(y + tz^0) \leq -\frac{1}{4}\|y\|_k^2 + \frac{3}{4}t^2 - C_\varepsilon C|t|^p$$

(here we have used again the Sobolev embedding). Note that all the constants here are independent on k . The last inequality implies that we can choose $\rho > 0$ large enough such that $J_k \leq 0$ on M_0 . Moreover, $\sup_M J_k(u) \leq K := \max(3t^2/4 - C_\varepsilon C|t|^q)$ and K is independent of k . Applying Theorem 2, we obtain the result.

The case ‘-’ sign we need only to apply Theorem 3 to the functional $-J_k$. \square

Remark 2 If we drop assumption (vi), i.e. allow 0 to be in the spectrum $\sigma(L)$, we still can use linking argument to prove the existence of k -periodic solutions. This can be done as in the case of the Dirichlet problem in a bounded domain (see [38], Section 2.4). However, in this case we have no uniform (with respect to k) bounds for the solution.

Remark 3 In the case ‘-’, if we drop the requirement $\inf \sigma(L) < 0$ then there is no nontrivial solution in E_k , as well as in E . This follows from the obvious fact that in this case the functionals $-J_k$ and $-J$ have strict global maximum at the origin.

3 Passage to the limit and finite action solutions

Now we are going to pass to the limit as $k \rightarrow \infty$ to obtain a finite action solution $u \in E$. The principal difficulty is to show that the limit is a non-zero function. With this aim we need some concentration compactness argument.

Lemma 3 *Let $Q_{(n)}$ be the cube with the edge length $l_n \rightarrow \infty$ centered at the origin and $K_r(\xi)$ the cube centered at ξ with the edge length r . Let $u_{(n)} \in H_{\text{loc}}^1(\mathbb{R}^N)$ be a sequence of l_n -periodic functions such that $\|u_{(n)}\|_{H^1(Q_{(n)})} \leq C$. Assume that there exists $r > 0$ such that*

$$\lim_{n \rightarrow \infty} \left(\sup_{\xi} \int_{K_r(\xi)} |u_{(n)}|^2 dx \right) = 0.$$

Then $\|u_{(n)}\|_{L^s(Q_{(n)})} \rightarrow 0$ for all $s \in (2, 2^)$.*

Proof. Denote by $Q'_{(n)}$ and $Q''_{(n)}$ the cubes with the sizes $l_n + r$ and $l_n + 2r$, respectively, centered at 0. Choose a cut-off function $\chi_n \in C_0^\infty(\mathbb{R}^N)$ such that $\text{supp } \chi_n \subset Q''_{(n)}$, $0 \leq \chi_n(x) \leq 1$, $\chi_n = 1$ on $Q'_{(n)}$ and $|\nabla \chi_n(x)| \leq C$ where $C > 0$ does not depend on n . Evidently, such a function exists.

Let us set $v_n = \chi_n u_{(n)}$. For n large enough, it is easy to verify, using the periodicity of $u_{(n)}$, that

$$\sup_{\xi} \int_{K_r(\xi)} |v_n|^2 dx = \sup_{\xi} \int_{K_r(\xi)} |u_{(n)}|^2 dx$$

and the sequence v_n is bounded in $H^1(\mathbb{R}^N)$. By the well-known lemma of P. L. Lions (see [20], Lemma I.1, and [38], Lemma 1.21), $v_n \rightarrow 0$ in $L^s(\mathbb{R}^N)$, $s \in (2, 2^*)$. Since $v_n = u_{(n)}$ on $Q_{(n)}$, we obtain the required. \square

Lemma 4 *Let $u_k \in E_k$ be a sequence such that $\|u_k\|_k$ is bounded and $J'_k(u_k) \rightarrow 0$. Then, passing to a subsequence still labelled by k , either*

$$1^0 \quad \|u_k\|_k \rightarrow 0 \text{ as } k \rightarrow \infty$$

or

2^0 *there exist a sequence of points $\xi_k \in \mathbb{R}^N$ and positive numbers r and η such that*

$$\lim_{k \rightarrow \infty} \int_{K_r(\xi_k)} |u_k|^2 dx \geq \eta.$$

Proof. Assume that 2^0 does not hold (along a subsequence still labelled by k). Since $J'_k(u_k) \rightarrow 0$, there exists a sequence $\varepsilon_k \rightarrow 0$ such that $\langle J'_k(u_k), v \rangle \leq \varepsilon_k \|v\|_k$ for any $v \in E_k$. Taking $v = P_k^+ u_k$ and $v = P_k^- u_k$ we obtain, as in the proof of Lemma 2,

$$\|u_k\|_k^2 \leq \varepsilon \|u_k\|_k^2 + C_\varepsilon \|u_k\|_{L^p(Q_k)}^p + C\varepsilon_k \|u_k\|_k.$$

By Lemma 3, $\|u_k\|_{L^p(Q_k)}^p \rightarrow 0$ and we obtain the required. \square

Theorem 5 *Let $u_k \in E_k$ be a sequence of nontrivial critical points of J_k such that the sequence $c_k = J_k(u_k)$ is bounded. Then there exists a nontrivial solution $u \in E = H^1(\mathbb{R}^N)$ of equation (1). Moreover, there exist a sequence $b_k \in \mathbb{Z}^N$ such that $u_k(x + b_k)$ converges to u in $H_{\text{loc}}^1(\mathbb{R}^N)$, up to the passage to a subsequence.*

Proof. By Lemma 1, the norms $\|u_k\|_k$ are bounded. Lemma 2 implies that $\|u_k\| - k \geq \varepsilon_1 > 0$. Therefore, case 1^o of Lemma 4 is not possible. From Lemma 4, 2^o, we obtain that, along a subsequence,

$$\|u_k\|_{L^2(K_r(\xi_k))}^2 \geq \eta/2.$$

There exists a sequence of integer vectors b_k such that the sequence of cubes $K_r(\xi_k - b_k)$ is confined in a bounded region. Set $\tilde{u}_k(x) = u_k(x + b_k)$. We have

$$\|u_k\|_{L^2(Q_{k_0})}^2 \geq \eta/2 \tag{7}$$

for $k \geq k_0$ and some k_0 . Since V and f are \mathbb{Z}^N -periodic, \tilde{u}_k are critical points of J_k and the norms $\|\tilde{u}_k\|_k$ are bounded. Passing to a subsequence, one can assume that the sequence \tilde{u}_k converges weakly in $H_{\text{loc}}^1(\mathbb{R}^N)$ to a function u . Since $\|\tilde{u}_k\|_k$ is bounded, $u \in H^1(\mathbb{R}^N)$. By the compactness of Sobolev embedding, $\tilde{u}_k \rightarrow u$ in $L_{\text{loc}}^s(\mathbb{R}^N)$, $2 \leq s < 2^*$, and, hence, $f(x, \tilde{u}_k) \rightarrow f(x, u)$ in $L_{\text{loc}}^1(\mathbb{R}^N)$. Now for any test function $\varphi \in C_0^\infty$ we have

$$\begin{aligned} \langle J'(u), \varphi \rangle &= \int_{\mathbb{R}^N} (\nabla u \nabla \varphi + V u \varphi - f(x, u) \varphi) dx = \\ &= \lim_{k \rightarrow \infty} \int_{\mathbb{R}^N} (\nabla \tilde{u}_k \nabla \varphi + V \tilde{u}_k \varphi - f(x, \tilde{u}_k) \varphi) dx = \\ &= 0. \end{aligned}$$

Therefore, u is a weak solution of (1) and $u \neq 0$ in view of (7).

Now let us prove the last statement. Let $\chi \in C_0^\infty(\mathbb{R}^N)$ and $\text{supp } \chi$ is contained in an open bounded set Ω . It is easy to verify the following identity

$$L(\chi \tilde{u}_k - \chi u) = h_k^1 + h_k^2 + h_k^3$$

where

$$\begin{aligned} h_k^1 &= \chi[f(x, \tilde{u}_k) - f(x, u)], \\ h_k^2 &= -\nabla \chi \nabla(\tilde{u}_k - u), \\ h_k^3 &= -(\Delta \chi)(\tilde{u}_k - u). \end{aligned}$$

We see that $h_k^1 \rightarrow 0$ in $L^{p-1}(\Omega)$, $h_k^2 \rightarrow 0$ weakly in $L^2(\Omega)$ and $h_k^3 \rightarrow 0$ in $L^2(\Omega)$. Since the space $L^s(\Omega)$, $2 \leq s < 2^*$, is compactly embedded into $H^{-1}(\mathbb{R}^N)$, then $h_k^1 + h_k^2 + h_k^3 \rightarrow 0$ in the last space. Assumption (vi) implies that the operator L considered as a bounded linear operator from $H^1(\mathbb{R}^N)$ into $H^{-1}(\mathbb{R}^N)$ has a bounded inverse operator. Therefore, $(\chi \tilde{u}_k - \chi u) \rightarrow 0$ in $H^1(\mathbb{R}^N)$ and we are done. \square

Combining Theorems 4 and 5, we obtain

Theorem 6 *Equation (1) possesses a nontrivial solution in $H^1(\mathbb{R}^N)$.*

4 Ground states

We say that a nontrivial solution $u \in E$ (resp., $u \in E_k$) of equation (1) is a *ground state* (resp., a *periodic ground state*) if the corresponding critical value of the functional $\pm J$ (resp., $\pm J_k$) is minimal possible among all nontrivial critical values of that functional. Since the functional $\pm J_k$ satisfies the Palais-Smale condition, the existence of periodic ground states follows immediately from Theorem 4 and Lemma 2. The case of ground states is more delicate. To treat it we need the following

Theorem 7 *In addition to assumptions (i) – (vi), suppose that*

$$|f(x, u + v) - f(x, u)| \leq C|v|(1 + |u|^{p-1}) \quad \text{if } |v| \leq \varepsilon \quad (8)$$

with some positive C and ε .

(a) *Let $u_k \in E$ be a bounded sequence such that $J'(u_k) \rightarrow 0$ and $J(u_k) \rightarrow c > 0$. Then there exist critical points u^i of J and sequences $b_k^i \in \mathbb{Z}^N$, $i = 1, \dots, n$ such that*

$$\sum_{i=1}^n J(u^i) = c \quad (9)$$

and

$$\|u_k - \sum_{i=1}^n u^i(\cdot + b_k^i)\| \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

(b) *Let $u_k \in E_k$ be a sequence such that $\|u_k\|_k$ is bounded, $J'_k(u_k) \rightarrow 0$ and $J_k(u_k) \rightarrow c > 0$. Then there exist critical points u^i of J satisfying (9) and sequences $b_k^i \in \mathbb{Z}^N$, $i = 1, \dots, n$ such that*

$$\|u_k - \sum_{i=1}^n u^i(\cdot + b_k^i)\|_{H^1(Q_k)} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

For the proof of part (a) we refer to [17]. Part (b) is borrowed from [25].

From Theorem 7 and previous remarks it follows

Theorem 8 *Under assumptions (i) – (vi) and (8) there exist at least one k -periodic ground state and at least one ground state.*

Now we want to study the behavior of k -periodic ground states as $k \rightarrow \infty$. We replace assumption (v) by the following stronger assumption

(v') *The function $f(x, u)$ is C^1 in u , f'_u is a Carathéodory function and there exists $\theta \in (0, 1)$ such that*

$$0 < u^{-1}f(x, u) \leq \theta f'_u(x, u)$$

for every $u \neq 0$ and

$$|f'_u(x, u)| \leq C(1 + |u|)^{p-2} \quad (10)$$

for all $u \in \mathbb{R}$, with $C > 0$.

This implies, in particular, that

$$0 < F(x, u) \leq \frac{\theta}{1 + \theta} f(x, u)u \quad (11)$$

for $u \neq 0$, and $\theta/(1 + \theta) < 1/2$. Therefore, (v) follows from (v'). Moreover, in this case inequality (8) holds as well. The functionals J_k and J are C^2 , and

$$\begin{aligned} \langle J_k''(u)w, v \rangle &= (P_k^+ w, v) - (P_k^- w, v) \pm \int_{Q_k} f'(x, u)wv \, dx = \\ &= \int_{Q_k} (\nabla w \nabla v + V(x)wv \pm f'(x, u)wv) \, dx, \end{aligned} \quad (12)$$

$$\begin{aligned} \langle J''(u)w, v \rangle &= (P^+ w, v) - (P^- w, v) \pm \int_{\mathbb{R}^N} f'(x, u)wv \, dx = \\ &= \int_{\mathbb{R}^N} (\nabla w \nabla v + V(x)wv \pm f'(x, u)wv) \, dx. \end{aligned} \quad (13)$$

Define the set $S_k^\pm \subset E_k$ that consists of all nonzero $u \in E_k$ such that

$$I_k(u) := \langle J_k'(u), u \rangle = 0$$

and

$$\langle J_k'(u), v \rangle = 0 \quad \forall v \in E_k^\mp.$$

Similarly, the set $S^\pm \subset E$ consists of all nonzero $u \in E$ such that

$$I(u) := \langle J'(u), u \rangle = 0$$

and

$$\langle J'(u), v \rangle = 0 \quad \forall v \in E^\mp.$$

These sets are nonempty, because they contain solutions.

Now let us consider the following minimization problems

$$m_k^\pm = \inf\{\pm J_k(u) : u \in S_k^\pm\} \quad (14)$$

and

$$m^\pm = \inf\{\pm J(u) : u \in S^\pm\}. \quad (15)$$

Inspection of the proof of Lemmas 1 and 2 gives us the following result.

Lemma 5 *There exist constants $\varepsilon_1 > 0$ and $\varepsilon_2 > 0$ independent on k such that for every $u_k \in S_k^\pm$ and every $u \in S^\pm$ we have $\|u_k\|_k \geq \varepsilon_1$, $\|u\| \geq \varepsilon_1$, $\pm J_k(u_k) \geq \varepsilon_2$ and $\pm J(u) \geq \varepsilon_2$. Moreover,*

$$\|u_k\|_k \leq C(|J_k(u_k)|^{1/2} + |J_k(u_k)|^{1/p'}), \quad \|u\| \leq C(|J(u)|^{1/2} + |J(u)|^{1/p'}).$$

Let $\overline{E}_k^\pm = \mathbb{R} \oplus E_k^\pm$ and $\overline{E}^\pm = \mathbb{R} \oplus E^\pm$. Consider the operators $G_k^\pm : E \rightarrow \overline{E}_k^\mp$ and $G^\pm : E \rightarrow \overline{E}^\mp$ defined by

$$G_k^\pm(u) = (\langle J'_k(u), u \rangle, P_k^\mp J'_k(u)),$$

$$G^\pm(u) = (\langle J'(u), u \rangle, P^\mp J'(u)),$$

respectively. Obviously, $S_k^\pm = (G_k^\pm)^{-1}(0) \setminus \{0\}$ and $S^\pm = (G^\pm)^{-1}(0) \setminus \{0\}$. Moreover, these operators are C^1 , and

$$(G_k^\pm)'(u)v = (\langle J''_k(u)v, u \rangle + \langle J'_k(u), v \rangle, P_k^\mp J''_k(u)v), \quad (16)$$

$$(G^\pm)'(u)v = (\langle J''(u)v, u \rangle + \langle J'(u), v \rangle, P^\mp J''(u)v). \quad (17)$$

Lemma 6 *The set S_k^\pm (respectively, S^\pm) is a closed C^1 -submanifold of E_k (respectively, E).*

Proof. We consider S^+ only. The remaining cases are similar. The result follows from the implicit function theorem if we check that G' is onto at every point of S^+ . For notational convenience we skip the superscript in G and the domain of integration.

Let $u_0 \in E$ ($u_0 \neq 0$), $\langle J'(u_0), u_0 \rangle = \alpha_0$ and $P^- \nabla J(u_0) = \alpha$. Identifying \overline{E}^- and the subspace $\mathbb{R}u_0 \oplus E^-$ via $(\tau, h) = \tau u_0 + h$, a direct calculation gives

$$\begin{aligned} (G'(u_0)(\tau, h), (\tau, h)) &= 2\alpha_0\tau^2 - 3\tau(P^-u_0, h) + \tau^2 \int [f(x, u_0)u_0 - f'_u(x, u_0)u_0^2] dx - \\ &\quad - (h, h) - \tau \int [2f'_u(x, u_0)u_0 + f(x, u_0)]h dx - \int f'_u(x, u_0)h^2 dx. \end{aligned}$$

Since for $h \in E^-$

$$\begin{aligned} (P^+u_0, h) - (P^-u_0, h) - \int f(x, u_0)h dx &= \\ = -(P^-u_0, h) - \int f(x, u_0)h dx &= (\alpha, h), \end{aligned}$$

we obtain

$$\begin{aligned} (G'(u_0)(\tau, h), (\tau, h)) &= 2\alpha_0\tau^2 - (h, h) + 3\tau(\alpha, h) - \\ - \int \{[f'_u(x, u_0)u_0^2 - f(x, u_0)u_0]\tau^2 &+ 2[f'_u(x, u_0)u_0 - f(x, u_0)]\tau h + f'_u(x, u_0)h^2\} dx = \\ = 2\alpha_0\tau^2 - (h, h) + 3\tau(\alpha, h) - \int (A\tau^2 &+ 2C\tau h + Bh^2) dx. \end{aligned}$$

Note that $B \geq 0$, and $B(x) = 0$ iff $u_0(x) = 0$. Hence, in the last case also $A(x) = C(x) = 0$. Since

$$A\tau^2 + 2C\tau h + Bh^2 = \left(A - \frac{C^2}{B}\right)\tau^2 + \left(\sqrt{B}h + \frac{C\tau}{\sqrt{B}}\right),$$

$$|\tau(\alpha, h)| \leq \frac{1}{2} \|\alpha\| (\tau^2 + \|h\|^2)$$

and, due to assumption (v'),

$$\begin{aligned} A - \frac{C^2}{B} &= \left(u_0 - \frac{f(x, u_0)}{f'_u(x, u_0)} \right) f(x, u_0) \geq \\ &\geq (1 - \theta) f(x, u_0) u_0, \end{aligned}$$

we obtain

$$\begin{aligned} (G'(u_0)(\tau, h), (\tau, h)) &\leq 2\alpha_0 \tau^2 - \|h\|^2 + \frac{3}{2} \|\alpha\| \tau^2 + \frac{3}{2} \|\alpha\| \|h\|^2 - \\ &\quad - \tau^2 (1 - \theta) \int f(x, u_0) u_0 dx. \end{aligned} \tag{18}$$

Now if $u_0 \in S^+$, then $\alpha_0 = 0$ and $\alpha = 0$. Equation (16) implies that on the subspace $\mathbb{R}u_0 \oplus E^- \subset E$ the operator $G'(u_0)$ is strictly negative defined, hence, invertible. Therefore, the implicit function theorem applies and we conclude. \square

Remark 4 It is clear that S_k^+ has a finite co-dimension, while S_k^- is finite dimensional. The tangent space to S_k^\pm at u_0 consists of all $h \in E_k$ such that

$$\langle I'_k(u_0), h \rangle = 0, \quad P_k^\mp \nabla J_k(u_0) h = 0$$

and similarly for S^\pm . The subspace $\mathbb{R}u_0 \oplus E_k^\mp \subset E_k$ (resp., $\mathbb{R}u_0 \oplus E^\mp \subset E$) is transverse to S_k^\pm (resp., to S^\pm) at u_0 , as it follows from the proof of Lemma 6.

Lemma 7 *Any critical point of the restriction of J_k to S_k (respectively, J to S) is a critical point of J_k (respectively, J). In particular, solutions of problems (14) and (15) are periodic ground states and ground states, respectively.*

Proof. A direct calculation shows that if $u_0 \in S_k^\pm$ (resp., $u_0 \in S^\pm$), then $J'_k(u_0)$ (resp., $J'_k(u_0)$) vanishes on the subspace $\mathbb{R}u_0 \oplus E_k^\mp \subset E_k$ (resp., $\mathbb{R}u_0 \oplus E^\mp \subset E$). Therefore, if u_0 is a critical point of J_k (resp., J) restricted to S_k^\pm (resp., to S^\pm), then $J'_k(u_0)$ (resp., $J'_k(u_0)$) vanishes everywhere, since it is equal to 0 also on the tangent space. \square

Theorem 9 *Problem (14) has at least one solution which is a periodic ground state. Moreover, $m_k^\pm \geq \varepsilon_2$, with ε_2 from Lemma 5.*

Proof. We only sketch the existence of the minimization problem on S_k^+ . On this set we have

$$J_k(u) = J_k(u) - \frac{1}{2} \langle J'_k(u), u \rangle = \int_{Q_k} \left(\frac{1}{2} f(x, u) u - F(x, u) \right) dx. \tag{19}$$

Hence, due to assumption (v), $J_k(u)$ is bounded below on S_k^+ . Consider a minimization sequence $u_n \in S_k^+$. As it follows from the Ekeland principle (see, e.g., [40]), we can assume

that u_n is, in addition, a Palais-Smale sequence for $J_k|_{S_k^+}$. (In [40] functionals defined on a Banach space are considered, but arguments work in the case of functionals on C^1 Banach manifolds, with only minor change). We shall show that, in fact, u_n is a Palais-Smale sequence for the whole functional J_k . This is enough to pass to the limit and get the result.

Let $g_n = \nabla J_k(u_n)$ and g_n^t its component tangent to S_k^+ , i.e. the orthogonal projection of g_n onto the tangent space at u_n . Then $g_n^t \rightarrow 0$. We have to show that $g_n \rightarrow 0$. Inequality (18), with $u_0 = u_n$, $\alpha_0 = 0$ and $\alpha = 0$, implies that the operator $(G_k^+)'(u_n)$ has a right inverse operator, A_k^n . The image of this operator is the subspace $\overline{E}^-(u_n) = \mathbb{R}u_n \oplus E^-$ and its norm is bounded above by a constant $C > 0$ independently of n . The operator $A_k^n(G_k^+)'(u_n)$ is the projector, P_k^n , onto $\overline{E}^-(u_n)$ parallel to the tangent space at u_n . The adjoint operator $(P_k^n)^*$ is the projector onto the orthogonal complement to $\overline{E}^-(u_n)$ parallel to the normal subspace at u_n . Hence, $g_n = (P_k^n)^*g_n^t$ and $\|g_n\|_k \leq C\|g_n^t\|$. This completes the proof. \square

Now we impose the following additional assumption

(vii) *There exist $C > 0$ and $\gamma \in (0, 1]$ such that*

$$|f'_u(x, u_1) - f'_u(x, u_2)| \leq C(1 + |u_1| + |u_2|)^{p-2-\gamma}|u_1 - u_2|^\gamma$$

for all $u_1, u_2 \in \mathbb{R}$.

This assumption implies, in particular, (10).

Theorem 10 *Under assumptions (i)–(iv), (v'), (vi) and (vii) we have*

$$m^\pm = \lim_{k \rightarrow \infty} m_k^\pm. \quad (20)$$

Moreover, let $u_k \in E_k$ be a solution of (14). Then, after passage to a subsequence still denoted by u_k , there exist a solution u of (15) and a sequence $b_k \in \mathbb{Z}^N$ such that

$$\|u_k - u(\cdot + b_k)\|_{H^1(Q_k)} \rightarrow 0. \quad (21)$$

Proof. We again consider only the case of “+” sign. Since u_1 is also a k -periodic solution, we see that $m_k^+ \leq m_1^+$. Hence, the sequence m_k^+ is bounded and Lemma 5 shows that the sequence $\|u_k\|_k$ is bounded. Due to Theorem 5 we can assume that u_k converges in $H_{\text{loc}}^1(\mathbb{R}^N)$ to a nontrivial solution $u \in E$. Note that $J(u) \geq m^+$.

Equation (19) implies that

$$m_k^+ = \int_{Q_k} \left(\frac{1}{2} f(x, u_k) u_k - F(x, u_k) \right) dx =: \int_{Q_k} g(x, u_k) dx.$$

Since the integrand here is nonnegative, we see that for any bounded domain $\Omega \subset \mathbb{R}^N$ and k large enough

$$m_k^+ \geq \int_{\Omega} g(x, u_k) dx$$

and, therefore,

$$\liminf m_k^+ \geq \int_{\Omega} g(x, u_k) dx .$$

Since Ω is an arbitrary domain and u is a nontrivial solution, we obtain

$$\liminf m_k^+ \geq \int_{\mathbb{R}^N} g(x, u_k) dx = J(u) \geq m^+ . \quad (22)$$

To prove (20), we have now to show that

$$\limsup m_k^+ \leq m^+ . \quad (23)$$

Let $v \in S^+$. Since $C_0^\infty(\mathbb{R}^N)$ is dense in E there exists a sequence of smooth functions $v_k \rightarrow v$ in E such that v_k has a compact support in Q_k . Having in mind periodic extension, we can also consider v_k as an element of E_k . Let $\alpha^k = P_k^- \nabla J_k(v_k)$ and $\alpha_0^k = I_k(v_k)$. Inequality (18), with $u_0 = v_k$, shows that the derivative of the map G_k^+ at v_k has a right inverse whose norm is bounded independently of k , provided k is large enough. Note that, due to assumption (vii), G_k^+ is Hölder equicontinuous, with the exponent γ , say, on the ball of radius $2\|v\|$ in E_k centered at the origin. Inspecting the proof of the implicit function theorem (see, e.g., [40]), we get right inverse map, T_k , to G_k^+ defined on the ball $B_k \subset \mathbb{R} \oplus E_k^-$ of radius r centered at (α_0^k, α^k) , where r is independent of k . Moreover, T_k is Hölder equicontinuous. For k large enough, $0 \in B_k$ and is close to the point (α_0^k, α^k) . Hence, $w_k = T_k(0)$ is close to v_k , $w_k \in S_k^+$ and

$$J_k(w_k) \leq J(v) + \varepsilon$$

if k is large enough. This implies (23).

The remaining part of the theorem follows from Theorem 7, part (b). \square

5 Exponential decay

Now let us study the decay of finite action solutions to equation (1).

Theorem 11 *Let $u \in H^1(\mathbb{R}^N)$ be a solution of (1). Then u is a continuous function and there exist positive constants C and λ such that*

$$|u(x)| \leq C \exp(-\lambda|x|)$$

where $\lambda \leq c \operatorname{dist}(0, \sigma(L)) = c\alpha$.

Proof. First, we have to show that $u \in L^\infty(\mathbb{R}^N)$. This can be done exactly as in the proof of Lemma 5.1, [8]. (That proof is based on the Sobolev estimates for Schrödinger operators). Next, Theorem B.3.3 of [35] implies that u is continuous. Now we set $W(x) := -f(x, u(x))/u(x)$ (with $W(x) = 0$ if $u(x) = 0$). Hence, u solves the equation

$$-\Delta + (V(x) + W(x))u = 0 \quad (24)$$

on \mathbb{R}^N . Since $V + W \in L^\infty(\mathbb{R}^N)$, Theorem C.3.1 of [35] shows that $u(x) \rightarrow 0$ as $x \rightarrow \infty$.

Since $W(x) \rightarrow 0$ as $x \rightarrow \infty$, the potential W defines a relatively compact perturbation of the operator L . Hence, $L + W$ has the same essential spectrum as L and may have only isolated eigenvalues of finite multiplicity outside the essential spectrum. Equation (24) means that 0 is an eigenvalue of $L + W$ outside the essential spectrum, hence, of finite multiplicity, and u is the corresponding eigenfunction. Now Theorem C.3.5, [35], provides the required exponential bound. \square

Remark 5 Using estimates for Green's function of L (see [18], Section 7.6.3), one can improve the estimate for λ as follows: $a \leq c\alpha^{1/2}$.

6 Further results

Now we discuss assumption (vi). More precisely, what may happen if 0 is in the spectrum of L ? It is commonly believed that in this case well localized solutions of (1) do not exist. As we see from equation (24), the question is closely related to the problem of absence of embedded eigenvalues for periodic Schrödinger operators perturbed by a decaying potential. Unfortunately, there is no satisfactory general result on embedded eigenvalues, except the case $N = 1$. In the case $N = 1$ F. S. Rofe-Beketov [33] (see also [34]) has proved that $L + W$ has no embedded eigenvalues if $(1 + |x|)W(x) \in L^1(\mathbb{R})$. This implies immediately the following result.

Theorem 12 *Let $N = 1$. Suppose that assumptions (i) and (ii) are satisfied and, for some $r > 1$, $|f(x, u)| \leq c|u|^r$ for a.e. $x \in \mathbb{R}$ and u in a neighborhood of 0. Then equation (1) has no nontrivial weak solutions such that $(1 + |x|)|u(x)|^{r-1} \in L^1(\mathbb{R})$. In particular, there is no nontrivial exponentially decaying solutions.*

On the other hand, T. Bartsch and Y. Ding have found the following existence result.

Theorem 13 *In addition to (i)–(v), assume that V and g are continuous functions of all their arguments, there exist constants $r \in [q, 2^*)$, $s \in [p, 2^*)$, $a_1 > 0$ and $a_2 > 0$ such that*

$$a_1|u|^r \leq F(x, u),$$

$$|f(x, u)| \leq a_2(|u|^{p-1} + |u|^{s-1})$$

for all $x \in \mathbb{R}^N$ and $u \in \mathbb{R}$. Let $0 \in \sigma(L)$ and there exists $b > 0$ such that $(0, b) \cap \sigma(L) = \emptyset$ in the case of '+' sign in (1) and $(-b, 0) \cap \sigma(L) = \emptyset$ in the case of '-' sign. Then equation (1) has a nontrivial weak solution $u \in H_{\text{loc}}^2(\mathbb{R}^N)$. This solution lies in $L^t(\mathbb{R}^N)$ for $r \leq t \leq 2^$, is continuous and $u(x) \rightarrow 0$ as $x \rightarrow \infty$.*

Certainly, the solution obtained in the last theorem should be not well decaying. If $N = 1$, Theorem 12 implies that this solution cannot decay too fast.

Thus, we conjecture that if $0 \in \sigma(L)$, then nontrivial well-decaying at infinity solutions to periodic NLS 1 do not exist. To support this conjecture assume that $0 \in \sigma(L)$ and

is not a lower (resp., upper) edge of a spectral gap in the case of ‘+’ (resp., ‘-’) sign in (1). Let $k_j \rightarrow \text{infity}$ be a sequence of integers such that $0 \notin \sigma(L_{k_j})$. Recall that $\sigma(L_k)$ is discrete and the countable set $\cup \sigma(L_k)$ is dense in $\sigma(L)$. Therefore, k_j is a generic sequence. Let $u_j = u_{k_j} \in E_{k_j}$ be the k_j -periodic solution obtained by linking (see Remark 2) and $c_j = c_{k_j}$ be the corresponding linking critical value defined in Theorem 4. Note that it does not depend on particular choice of z^0 . Denote by α_j the distance from 0 to $\sigma(L_{k_j})$. Then we have

Proposition 1 *Under the assumptions imposed above*

$$\|u_j\|_{H^1(Q_{k_j})} \leq C\alpha_j^{\frac{q}{2(q-2)}},$$

where $C > 0$ does not depend on j .

Proof. Since the result seems to be not exact, we only sketch the proof in the case of ‘+’ sign.

Following the proof of Lemma 1 (see also Remark 1), we obtain

$$\|u_j\|_{H^1(Q_{k_j})} \leq C\alpha_j^{-1/2}(|c_j|^{1/p'} + |c_j|^{1/2}), \quad (25)$$

where p' is the conjugate exponent to p .

Now we estimate the linking critical value as in the proof of Theorem 4. We refine the argument after inequality (6). Take as $z^0 \in E_{k_j}^+$, the eigenvector of L_{k_j} with $\|z^0\|_{k_j} = 1$ that correspond to the first positive eigenvalue λ_0 of L_{k_j} . Then $\|z^0\|_{H^{-1}(Q_{k_j})} \leq C\|z^0\|_{L^q(Q_{k_j})}$, with $C > 0$ independent of j . Since

$$-\Delta z^0 + z^0 = \lambda_0 z^0 - V(x)z^0,$$

we have that $\|z^0\|_{H^1(Q_{k_j})} \leq C\|z^0\|_{L^q(Q_{k_j})}$. From (6) we get

$$c_j \leq \sup_M J_{k_j}(u) \leq \max(3t^2 - C\alpha_j^{-q/2}t^p).$$

A direct calculation gives us

$$c_j \leq C\alpha_j^{\frac{q}{q-2}}.$$

Since $1 < p' < 2$, this together with (25) implies the required. \square

Remark 6 Certainly, the conclusion of Proposition 1 holds also for periodic ground states.

Construction of ground states in Section s4 shows that the ground critical value does not exceed the linking value, which is not necessary a critical value in this case. Therefore, arguing as in Proposition 1, we obtain the following result.

Proposition 2 *Let $u \in E$ be a ground state of equation (1). Under the assumptions of Theorem 8 we have the estimate*

$$\|u\|_{H^1(\mathbb{R}^N)} \leq C(\alpha_{\pm})^{\frac{q}{2(q-2)}},$$

where $(-\alpha_-, \alpha_+)$ is the spectral gap containing 0.

As consequence, we see that ground states bifurcate from the trivial solution corresponding to an appropriate edge of the spectral gap.

7 Gap solitons

In this section we give an application of the previous results to a problem that arises in the theory of photonic crystals, the existence of gap solitons. Photonic crystals are dielectric media with spatially periodic (or close-to-periodic) structure. A good introduction into this field can be found in [16]. For a survey of rigorous mathematical results we refer to [18]. Both these publications deal with linear optical media. If the medium we consider is nonlinear, many new phenomena occur. Among them one of the most interesting is the possibility of gap solitons, i.e. spatially localized light patterns with the frequency prohibited by the linear theory. On physical level a simple description of this phenomenon is presented in [22]. However, up to now there was no mathematically rigorous proof of the existence of gap solitons even in simplest situations. Here we show that some existence result can be extracted from the results on periodic NLS.

In a dielectric medium we consider the system of Maxwell equations

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t},$$

$$\nabla \times \mathbf{H} = \frac{\partial \mathbf{D}}{\partial t},$$

$$\nabla \mathbf{D} = 0,$$

$$\nabla \mathbf{B} = 0.$$

Since we assume that the medium is non-magnetic, we set $\mathbf{B} = \mathbf{H}$. We consider the following constitutive relation between displacement and electric fields:

$$\mathbf{D} = (\varepsilon(x) + \chi(x)\langle |\mathbf{E}|^2 \rangle)\mathbf{E}$$

where $\langle \cdot \rangle$ stands for time average. Such type of nonlinear response was introduced by N. N. Akhmediev [3] (see also [36] and references therein). The cubic form of the nonlinearity means that we concentrate on Kerr-like media.

In the case of two dimensional structure we assume that the functions $\varepsilon(x) = \varepsilon(x_1, x_2)$ and $\chi(x) = \chi(x_1, x_2)$ are independent on x_3 , periodic in (x_1, x_2) and of the class L^∞ . Moreover, we suppose that $\varepsilon(x) \geq \varepsilon_0 > 0$ and either $\chi(x) > 0$, or $\chi(x) < 0$ everywhere.

The assumption means that the medium we consider is either everywhere self-focusing, or everywhere defocusing [22]. We restrict ourself to the so-called E -mode $\mathbf{E} = (0, 0, E)$ and look for solutions of the form

$$E = u(x_1, x_2) \cos(\beta x_3 - \omega t + \theta_0) \quad (26)$$

where the amplitude vanishes at infinity. Such solutions represent light patterns of the frequency ω that are localized in (x_1, x_2) directions and propagate along the x_3 -axis, with the wave number β . Obviously, the problem reduces to the two dimensional NLS

$$-\Delta u - \omega^2 \varepsilon u + \beta^2 u = \omega^2 \chi u^3. \quad (27)$$

Now we can apply the previous results. In the self-focusing case ($\chi > 0$) we obtain a nontrivial exponentially decaying solution if $-\beta^2$ is not in the spectrum of the operator $L_\omega := -\Delta - \omega^2 \varepsilon$. In particular, given ω such a solution exists for all $|\beta|$ large enough. In the defocusing case ($\chi < 0$) nontrivial exponentially localized solutions exist, provided $-\beta^2$ is not in $\sigma(L_\omega)$, but *not* below the spectrum. This means that $-\beta^2$ must belong to a *finite* spectral gap of L_ω .

The case $\beta = 0$ is of particular interest. Let us assume that $0 \notin \sigma(L_\omega)$. This means that ω is a prohibited frequency [18]. Since the potential of L_ω is *negative*, 0 belongs to a finite gap. Therefore, independently of the sign of χ there exists a nontrivial solution of the form $u(x_1, x_2) \cos(-\omega t + \theta_0)$ with exponentially decaying amplitude u . Such solutions represent so-called *standing* gap solitons. Hence, we obtain the existence of gap solitons for every prohibited frequency. In fact, in this case ($0 \in \sigma(L_\omega)$) there exists a nontrivial exponentially localized solution of the form (26) for all β close enough to 0.

Exactly the same results hold for one dimensional structures. This case reduces to the one dimensional periodic NLS.

Now let us discuss the behavior of gap solitons with respect to ω . Let $(-\alpha_-, \alpha_+)$ be the spectral gap of L_ω containing 0 and (ω_-, ω_+) the corresponding gap of (nonnegative) frequencies. Note that if ω goes to ω_\pm , then α_{pm} goes to 0. Indeed, the spectrum of L_ω is the union of closed intervals

$$\left[\min_{\theta \in [0, 2\pi]} \lambda_j(\theta), \max_{\theta \in [0, 2\pi]} \lambda_j(\theta) \right],$$

where $\lambda_j(\theta)$ are the Bloch eigenvalues (see, e.g., [32]). Since the potential $-\omega^2 \varepsilon(x)$ decreases monotonically as ω increases, the comparison principle for eigenvalues (see, e.g., [10]) tells us that the eigenvalues $\lambda_j(\theta)$ decreases monotonically and continuously, and we are done. Now Proposition 2 shows that gap solitons bifurcate from zero solutions corresponding to $\omega = \omega_+$ in the self-focusing case and $\omega = \omega_-$ in the defocusing case.

We have considered here the case when $\chi(x)$ does not change sign. The case of sign changing $\chi(x)$ is not less important, but completely open. It corresponds to a mixture of self-focusing and defocusing optical materials.

Another important problem is the existence of gap solitons in periodic media with *saturation*. In this case the nonlinearity is asymptotically linear. In the context of applications to nonlinear optical wave guides (one dimensional nonperiodic problem) such nonlinearities were studied extensively by C. Stuart (see, in particular, [36]).

References

- [1] Ackerman N., *On a periodic Schrödinger equation with nonlocal superlinear part*, Preprint, Giessen University, 2002.
- [2] Adams, *Sobolev Spaces*, Acad. Press, 1975.
- [3] Akhmediev N. N., Novel class of nonlinear surface waves. Asymmetric modes in a symmetric layered structure, *Sov. Phys. JETP*, **56** (1982), 299–303.
- [4] Alama S. and Li Y. Y., Existence of solutions for semilinear elliptic equations with indefinite linear part, *J. Differ. Equat.*, **96** (1992), 89–115.
- [5] Alama S. and Li Y. Y., On “multibump” bound states for certain semilinear elliptic equations, *Indiana Univ. Math. J.*, **41** (1992), 983–1026.
- [6] Arioli G. and Szulkin A., *A semilinear Schrödinger equation in the presence of a magnetic field*, Preprint, Stockholm University, 2002.
- [7] Bartsch T. and Ding Y., On a nonlinear Schrödinger equation with periodic potential, *Math. Ann.*, **313** (1999), 15–37.
- [8] Bartsch T., Pankov A. and Wang Zh.-Q., Nonlinear Schrödinger equations with steep potential well, *Commun. Contemp. Math.*, **3** (2001), 549–569.
- [9] Benci V. and Rabinowitz P. H., Critical point theorems for indefinite functionals, *Inv. Math.*, **52** (1979), 241–273.
- [10] Berezin F. and Shubin M., *The Schrödinger Equation*, Kluwer, Dordrecht, 1991.
- [11] Buffoni B., Jeanjean L. and Stuart C., Existence of nontrivial solutions to a strongly indefinite semilinear equation, *Proc. Amer. Math. Soc.*, **119** (1993), 179–186.
- [12] Chabrowski J. and Yang J., On Schrödinger equation with periodic potential and critical Sobolev exponent, *Topol. Methods Nonlinear Anal.*, **12** (1998), 245–261.
- [13] Coti Zelati V. and Rabinowitz P. H., Homoclinic type solutions for a semilinear elliptic PDE on \mathbb{R}^n , *Commun. Pure Appl. Math.*, **45** (1992), 1217–1269.
- [14] Courteille Ph., Bagnato V. and Yukalov V., Bose-Einstein condensation of trapped atomic gases, *Laser Physics*, **11** (2001), 659–800.
- [15] Jeanjean L., Solutions in spectral gaps for nonlinear equations of Schrödinger type, *J. Differ. Equat.*, **112** (1994), 53–80.
- [16] Joannopoulos J. D., Maede R. D. and Winn J. N., *Photonic Crystals, Molding the Flow of Light*, Princeton Univ. Press, NJ, 1995.

- [17] Kryszewski W. and Szulkin A., Generalized linking theorem with an application to semilinear Schrödinger equation, *Adv. Differ. Equat.*, **3** (1998), 441–472.
- [18] Kuchment P., The mathematics of photonic crystals, *Mathem. Modeling in Opt. Sci.*, 207–272, SIAM, 2001.
- [19] Li G. and Szulkin A., An asymptotically periodic Schrödinger equation with indefinite linear part, *Commun. Contemp. Math.*, **4** (2002), 763–776.
- [20] Lions P. L., The concentration compactness principle in the calculus of variations. The locally compact case, II, *Ann. Inst. H. Poincaré, Anal. Non Lin.*, **1** (1984), 223–283.
- [21] Maz'ya V. G., *Sobolev Spaces*, Springer, 1985.
- [22] Mills D. L., *Nonlinear Optics. Basic Concepts*, Springer, 1998.
- [23] Pankov A., *Introduction to Spectral Theory of Schrödinger Operators* (Lecture Notes), Preprint, Giessen University, 2001.
- [24] Pankov A., Semilinear elliptic equations on \mathbb{R}^n with nonstabilizing coefficients, *Ukr. Math. J.*, **41** (1989), 1075–1078, transl. from *Ukr. Mat. Zh.*, **41** (1989), 1247–1251.
- [25] Pankov A. and Pflüger K., On a semilinear Schrödinger equation with periodic potential, *Nonlin. Analysis*, **33** (1998), 593–609.
- [26] Pankov A. and Pflüger K., Periodic and solitary traveling waves for the generalized Kadomtsev-Petviashvili equations, *Math. Meth. Appl. Sci.*, **22** (1999), 733–752.
- [27] Pankov A. and Pflüger K., On ground traveling waves for the generalized Kadomtsev-Petviashvili equations, *Math. Phys., Analysis, Geom.*, **3** (2000), 33–47.
- [28] Pankov A. and Pflüger K., Traveling waves in lattice dynamical systems, *Math. Meth. Appl. Sci.*, **23** (2000), 1223–1235.
- [29] Rabinowitz P. H., *Minimax Methods in Critical Point Theory with Applications to Differential Equations*, AMS Reg. Conf. Ser. Math. **65**, 1986.
- [30] Rabinowitz P. H., A note on semilinear elliptic equation on \mathbb{R}^n , *Nonlinear Analysis: A Tribute in Honour of G. Prodi, Quad. Scu. Norm. Super. Pisa*, (1991), 307–318.
- [31] Rabinowitz P. H., On a class of nonlinear Schrödinger equations, *Z. Angew. Math. Phys.*, **43** (1992), 270–291.
- [32] Reed M. and Simon B., *Methods of Modern Mathematical Physics, vol. IV*, Acad. Press, 1978.

- [33] Rofe-Beketov F. S., A test for finiteness of the number of discrete levels introduced into the gap of a continuous spectrum by perturbation of a periodic potential, *Soviet Math. Dokl.*, **5** (1964), 689–692.
- [34] Rofe-Beketov F. S., Khol'kin A. M., *Spectral Analysis of Differential Operators*, World Sci., Singapore, to appear.
- [35] Simon B., Schrödinger semigroups, *Bull. Amer. Math. Soc. (N. S.)*, **7** (1982), 447–526.
- [36] Stuart C., Guidance properties of nonlinear planar waveguides, *Arch. Rat. Mech. Anal.*, **125** (1993), 145–200.
- [37] Trostler C. and Willem M., Nontrivial solutions of a semilinear Schrödinger equation, *Commun. Part. Differ. Equat.*, **21** (1996), 1431–1449.
- [38] Willem M., *Minimax Methods*, Birkhäuser, Boston, 1996.
- [39] Willem M. and Zou W., *On a semilinear Dirichlet problem and a nonlinear Schrödinger equation with periodic potential*, Preprint, Univ. Catholique Louvain, 2000.
- [40] Zeidler E., *Applied Functional Analysis. Main Principles and their Applications*, Springer, New York, 1995.