

# ON THE SOLVABILITY CONDITIONS FOR A LINEARIZED CAHN-HILLIARD EQUATION

Vitaly Volpert<sup>1</sup>, Vitali Vougalter<sup>2</sup>

<sup>1</sup> Institute Camille Jordan, UMR 5208 CNRS, University Lyon 1  
Villeurbanne, 69622, France  
e-mail: volpert@math.univ-lyon1.fr

<sup>2</sup> Department of Mathematics and Applied Mathematics, University of Cape Town  
Private Bag, Rondebosch 7701, South Africa  
e-mail: Vitali.Vougalter@uct.ac.za

**Abstract:** We derive solvability conditions in  $H^4(\mathbb{R}^3)$  for a fourth order partial differential equation which is the linearized Cahn-Hilliard problem using the results obtained for a Schrödinger type operator without Fredholm property in our preceding work [15].

**AMS Subject Classification:** 35J10, 35P10, 31A30

**Key words:** solvability conditions, non Fredholm operators, Sobolev spaces

## 1. Introduction

Consider a binary mixture and denote by  $c$  its composition, that is the fraction of one of its two components. Then the evolution of the composition is described by the Cahn-Hilliard equation (see, e.g. [1], [10]):

$$\frac{\partial c}{\partial t} = M\Delta \left( \frac{d\phi}{dc} - K\Delta c \right), \quad (1.1)$$

where  $M$  and  $K$  are some constants and  $\phi$  is the free energy density. In the stationary case we obtain the equation

$$\Delta(\Delta c + F(c)) = 0, \quad (1.2)$$

where  $F(c) = -\frac{1}{K} \frac{d\phi}{dc}$ . The existence, stability and some properties of solutions of the Cahn-Hilliard equation have been studied extensively in recent years (see e.g. [3], [6], [10]). In this work we investigate solvability of a linear fourth order equation which can be considered as a linearized Cahn-Hilliard equation:

$$\Delta(\Delta u + V(x)u + au) = f(x). \quad (1.3)$$

Here the potential  $V(x)$  is a smooth function vanishing at infinity. The precise conditions on it will be specified below. The function  $f(x)$  belongs to the appropriate weighted Hölder space, which will imply its square integrability, and  $a$  is a nonnegative constant. We will study this equation in  $\mathbb{R}^3$ .

Let us recall that Fredholm solvability conditions affirm that an operator equation  $Lu = f$  is solvable if and only if the right-hand side is orthogonal to all solutions  $w$  of the homogeneous adjoint equation  $L^*w = 0$ . This fundamental result is true if the operator  $L$  satisfies the Fredholm property, that is its kernel has a finite dimension, its image is closed, the codimension of the image is also finite.

The operator

$$Lu = \Delta(\Delta u + V(x)u + au)$$

considered as acting from  $H^4(\mathbb{R}^3)$  into  $L^2(\mathbb{R}^3)$  (or in the corresponding Hölder spaces) does not satisfy the Fredholm property. Indeed, since  $V(x)$  vanishes at infinity, then the essential spectrum of this operator is the set of all complex  $\lambda$  for which the equation

$$\Delta(\Delta u + au) = \lambda u$$

has a nonzero bounded solution. Applying the Fourier transform, we obtain

$$\lambda = -\xi^2(a - \xi^2), \quad x \in \mathbb{R}.$$

Hence the essential spectrum contains the origin. Consequently, the operator does not satisfy the Fredholm property, and solvability conditions of equation (1.3) are not known. We will obtain solvability conditions for this equation using the method developed in our previous papers [15], [16], [17], [18]. This method is based on spectral decomposition of self-adjoint operators.

Obviously, the problem above can be conveniently rewritten in the equivalent form of the system of two second order equations

$$\begin{cases} -\Delta v = f(x), \\ -\Delta u - V(x)u - au = v(x) \end{cases} \quad (1.4)$$

in which the first one has an explicit solution due to the fast rate of decay of its right side stated in Assumption 3, namely

$$v_0(x) := \frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{f(y)}{|x-y|} dy \quad (1.5)$$

with properties established in Lemma A1 of the Appendix. Note that both equations of the system above involve second order differential operators without Fredholm property. Their essential spectra are  $\sigma_{ess}(-\Delta) = [0, \infty)$  and  $\sigma_{ess}(-\Delta - V(x) -$

$a) = [-a, \infty)$  for  $V(x) \rightarrow 0$  at infinity (see e.g. [8]), such that neither of the operators has a finite dimensional isolated kernel. Solvability conditions for operators of that kind have been studied extensively in recent works for a single Schrödinger type operator (see [15]), sums of second order differential operators (see [16]), the Laplacian operator with the drift term (see [17]). Non Fredholm operators arise as well while studying the existence and stability of stationary and travelling wave solutions of certain reaction-diffusion equations (see e.g. [5], [7], [14]). For the second equation in system (1.4) we introduce the corresponding homogeneous problem

$$-\Delta w - V(x)w - aw = 0. \quad (1.6)$$

We make the following technical assumptions on the scalar potential and the right side of equation (1.3). Note that the first one contains conditions on  $V(x)$  analogous to those stated in Assumption 1.1 of [15] (see also [16], [17]) with the slight difference that the precise rate of decay is assumed not a.e. as before but pointwise since in the present work the potential function is considered to be smooth.

**Assumption 1.** *The potential function  $V(x) : \mathbb{R}^3 \rightarrow \mathbb{R}$  satisfies the estimate*

$$|V(x)| \leq \frac{C}{1 + |x|^{3.5+\delta}}$$

with some  $\delta > 0$  and  $x = (x_1, x_2, x_3) \in \mathbb{R}^3$  such that

$$4^{\frac{1}{9}} \frac{9}{8} (4\pi)^{-\frac{2}{3}} \|V\|_{L^\infty(\mathbb{R}^3)}^{\frac{1}{9}} \|V\|_{L^{\frac{8}{3}}(\mathbb{R}^3)}^{\frac{8}{9}} < 1 \quad \text{and} \quad \sqrt{c_{HLS}} \|V\|_{L^{\frac{3}{2}}(\mathbb{R}^3)} < 4\pi.$$

Here and further down  $C$  stands for a finite positive constant and  $c_{HLS}$  given on p.98 of [11] is the constant in the Hardy-Littlewood-Sobolev inequality

$$\left| \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{f_1(x)f_1(y)}{|x-y|^2} dx dy \right| \leq c_{HLS} \|f_1\|_{L^{\frac{3}{2}}(\mathbb{R}^3)}^2, \quad f_1 \in L^{\frac{3}{2}}(\mathbb{R}^3).$$

Here and below the norm of a function  $f_1 \in L^p(\mathbb{R}^3)$ ,  $1 \leq p \leq \infty$  is denoted as  $\|f_1\|_{L^p(\mathbb{R}^3)}$ .

**Assumption 2.**  $\Delta V \in L^2(\mathbb{R}^3)$  and  $\nabla V \in L^\infty(\mathbb{R}^3)$ .

We will use the notation

$$(f_1(x), f_2(x))_{L^2(\mathbb{R}^3)} := \int_{\mathbb{R}^3} f_1(x) \bar{f}_2(x) dx,$$

with a slight abuse of notations when these functions are not square integrable, like for instance some of those used in the Assumption 3 below. Let us introduce the auxiliary quantity

$$\rho(x) := (1 + |x|^2)^{\frac{1}{2}}, \quad x = (x_1, x_2, x_3) \in \mathbb{R}^3 \quad (1.7)$$

and the space  $C_a^\mu(\mathbb{R}^3)$ , where  $a$  is a real number and  $0 < \mu < 1$  consisting of all functions  $u$  for which

$$u\rho^a \in C^\mu(\mathbb{R}^3).$$

Here  $C^\mu(\mathbb{R}^3)$  stands for the Hölder space such that the norm on  $C_a^\mu(\mathbb{R}^3)$  is defined as

$$\|u\|_{C_a^\mu(\mathbb{R}^3)} := \sup_{x \in \mathbb{R}^3} |\rho^a(x)u(x)| + \sup_{x, y \in \mathbb{R}^3} \frac{|\rho^a(x)u(x) - \rho^a(y)u(y)|}{|x - y|^\mu}.$$

Then the space of all functions for which

$$\partial^\alpha u \in C_{a+|\alpha|}^\mu(\mathbb{R}^3), \quad |\alpha| \leq l,$$

where  $l$  is a nonnegative integer is being denoted as  $C_a^{\mu+l}(\mathbb{R}^3)$ . Let  $P(s)$  be the set of polynomials of three variables of the order less or equal to  $s$  for  $s \geq 0$  and  $P(s)$  is empty when  $s < 0$ . We make the following assumption on the right side of the linearized Cahn-Hilliard problem.

**Assumption 3.** *Let  $f(x) \in C_{6+\varepsilon}^\mu(\mathbb{R}^3)$  for some  $\varepsilon > 0$  and the orthogonality relation*

$$(f(x), p(x))_{L^2(\mathbb{R}^3)} = 0 \tag{1.8}$$

*holds for any polynomial  $p(x) \in P(3)$  satisfying the equation  $\Delta p(x) = 0$ .*

**Remark.** *A good example of such polynomials of the third order is*

$$\frac{a}{2}x_1^3 + \frac{b}{2}x_1x_2^2 + \frac{c}{2}x_1x_3^2,$$

*where  $a, b$  and  $c$  are constants, such that  $3a + b + c = 0$ . The set of admissible  $p(x)$  includes also constants, linear functions of three variables and many more examples.*

By means of Lemma 2.3 of [15], under our Assumption 1 above on the potential function, the operator  $-\Delta - V(x) - a$  is self-adjoint and unitarily equivalent to  $-\Delta - a$  on  $L^2(\mathbb{R}^3)$  via the wave operators (see [9], [13])

$$\Omega^\pm := s - \lim_{t \rightarrow \mp\infty} e^{it(-\Delta - V)} e^{it\Delta}$$

with the limit understood in the strong  $L^2$  sense (see e.g. [12] p.34, [4] p.90). Therefore,  $-\Delta - V(x) - a$  on  $L^2(\mathbb{R}^3)$  has only the essential spectrum  $\sigma_{ess}(-\Delta - V(x) - a) = [-a, \infty)$ . Via the spectral theorem, its functions of the continuous spectrum satisfying

$$[-\Delta - V(x)]\varphi_k(x) = k^2\varphi_k(x), \quad k \in \mathbb{R}^3, \tag{1.9}$$

in the integral formulation the Lippmann-Schwinger equation for the perturbed plane waves (see e.g. [12] p.98)

$$\varphi_k(x) = \frac{e^{ikx}}{(2\pi)^{\frac{3}{2}}} + \frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{e^{i|k||x-y|}}{|x-y|} (V\varphi_k)(y) dy \quad (1.10)$$

and the orthogonality relations

$$(\varphi_k(x), \varphi_q(x))_{L^2(\mathbb{R}^3)} = \delta(k - q), \quad k, q \in \mathbb{R}^3 \quad (1.11)$$

form the complete system in  $L^2(\mathbb{R}^3)$ . We introduce the following auxiliary functional space (see also [17], [18])

$$\tilde{W}^{2,\infty}(\mathbb{R}^3) := \{w(x) : \mathbb{R}^3 \rightarrow \mathbb{C} \mid w, \nabla w, \Delta w \in L^\infty(\mathbb{R}^3)\}. \quad (1.12)$$

As distinct from the standard Sobolev space we require here not the boundedness of all second partial derivatives of the function but of its Laplacian. Our main result is as follows.

**Theorem 4.** *Let Assumptions 1, 2 and 3 hold,  $a \geq 0$  and  $v_0(x)$  is given by (1.5). Then problem (1.3) admits a unique solution  $u_a \in H^4(\mathbb{R}^3)$  if and only if*

$$(v_0(x), w(x))_{L^2(\mathbb{R}^3)} = 0 \quad (1.13)$$

for any  $w(x) \in \tilde{W}^{2,\infty}(\mathbb{R}^3)$  satisfying equation (1.6).

**Remark.** *Note that  $\varphi_k(x) \in \tilde{W}^{2,\infty}(\mathbb{R}^3)$ ,  $k \in \mathbb{R}^3$ , which was proven in Lemma A3 of [17]. By means of (1.9) these perturbed plane waves satisfy the homogeneous problem (1.6) when the wave vector  $k$  belongs to the sphere in three dimensions centered at the origin of radius  $\sqrt{a}$ .*

## 2. Proof of the main result

Armed with the technical lemma of the Appendix we proceed to prove the main result.

*Proof of Theorem 4.* The linearized Cahn-Hilliard equation (1.3) is equivalent to system (1.4) in which the first equation admits a solution  $v_0(x)$  given by (1.5). The function  $v_0(x) \in L^2(\mathbb{R}^3) \cap L^\infty(\mathbb{R}^3)$  and  $|x|v_0(x) \in L^1(\mathbb{R}^3)$  by means of Lemma A1 and Assumption 3. Then according to Theorem 3 of [18] the second equation in system (1.4) with  $v_0(x)$  in its right side admits a unique solution in  $H^2(\mathbb{R}^3)$  if and only if the orthogonality relation (1.13) holds. This solution of problem (1.3)  $u_a(x) \in H^2(\mathbb{R}^3) \subset L^\infty(\mathbb{R}^3)$  via the Sobolev embedding theorem,  $a \geq 0$  satisfies the equation

$$-\Delta u_a - V(x)u_a - au_a = v_0(x).$$

We use the formula

$$\Delta(Vu_a) = V\Delta u_a + 2\nabla V \cdot \nabla u_a + u_a\Delta V \quad (2.14)$$

with the “dot” denoting the standard scalar product of two vectors in three dimensions. The first term in the right side of (2.14) is square integrable since  $V(x)$  is bounded and  $\Delta u_a(x) \in L^2(\mathbb{R}^3)$ . Similarly  $u_a\Delta V \in L^2(\mathbb{R}^3)$  since  $u_a(x)$  is bounded and  $\Delta V$  is square integrable by means of Assumption 2. For the second term in the right side of (2.14) we have  $\nabla u_a(x) \in L^2(\mathbb{R}^3)$  and  $\nabla V$  is bounded via Assumption 2, which yields  $\nabla V \cdot \nabla u_a \in L^2(\mathbb{R}^3)$  and therefore,  $\Delta(Vu_a) \in L^2(\mathbb{R}^3)$ . The right side of problem (1.3) belongs to  $L^2(\mathbb{R}^3)$  due to Assumption 3. Indeed, since  $\sup_{x \in \mathbb{R}^3} |\rho^{6+\varepsilon} f| \leq C$ , we arrive at the estimate

$$|f(x)| \leq \frac{C}{(\rho(x))^{6+\varepsilon}}, \quad x \in \mathbb{R}^3 \quad (2.15)$$

with  $\rho(x)$  defined explicitly in (1.7). Hence from equation (1.3) we deduce that  $\Delta^2 u_a \in L^2(\mathbb{R}^3)$ . Any partial third derivative of  $u_a$  is also square integrable due to the trivial estimate in terms of the  $L^2(\mathbb{R}^3)$  norms of  $u_a$  and  $\Delta^2 u_a$ , which are finite. This implies that  $u_a \in H^4(\mathbb{R}^3)$ .

To investigate the issue of uniqueness we suppose  $u_1, u_2 \in H^4(\mathbb{R}^3)$  are two solutions of problem (1.3). Then their difference  $u(x) = u_1(x) - u_2(x) \in H^4(\mathbb{R}^3)$  satisfies equation (1.3) with vanishing right side. Clearly  $u, \Delta u \in L^2(\mathbb{R}^3)$  and  $Vu \in L^2(\mathbb{R}^3)$ . Therefore,  $v(x) = -\Delta u - V(x)u - au \in L^2(\mathbb{R}^3)$  and solves the equation  $\Delta v = 0$ . Since the Laplace operator does not have any nontrivial square integrable zero modes,  $v(x) = 0$  a.e. in  $\mathbb{R}^3$ . Hence, we arrive at the homogeneous problem  $(-\Delta - V(x) - a)u = 0$ ,  $u(x) \in L^2(\mathbb{R}^3)$ . The operator in brackets is unitarily equivalent to  $-\Delta - a$  on  $L^2(\mathbb{R}^3)$  as discussed above and therefore  $u(x) = 0$  a.e. in  $\mathbb{R}^3$ . ■

### 3. Appendix

**Lemma A1.** *Let Assumption 3 hold. Then  $v_0(x) \in L^2(\mathbb{R}^3) \cap L^\infty(\mathbb{R}^3)$  and  $xv_0(x) \in L^1(\mathbb{R}^3)$ .*

*Proof.* According to the result of [2], for the solution of the Poisson equation (1.5) under the condition  $f(x) \in C_{6+\varepsilon}^\mu(\mathbb{R}^3)$  and orthogonality relation (1.8) given in Assumption 3 we have  $v_0(x) \in C_{4+\varepsilon}^{\mu+2}(\mathbb{R}^3)$ . Hence  $\sup_{x \in \mathbb{R}^3} |\rho^{4+\varepsilon} v_0| \leq C$ , such that

$$|v_0(x)| \leq \frac{C}{(\rho(x))^{4+\varepsilon}}, \quad x \in \mathbb{R}^3.$$

The statement of the lemma easily follows from definition (1.7). ■

**Remark.** *Note that the boundedness of  $v_0(x)$  can be easily shown via the argument of Lemma 2.1 of [15], which relies on Young’s inequality. The square*

integrability of  $v_0(x)$  can be proven by applying the Fourier transform to it, using the facts that  $f(x) \in L^2(\mathbb{R}^3)$ ,  $|x|f(x) \in L^1(\mathbb{R}^3)$  and its Fourier image vanishes at the origin since it is orthogonal to a constant by means of Assumption 3.

## References

- [1] N.D. Alikakos, G. Fusco, *Slow dynamics for the Cahn-Hilliard equation in higher space dimensions. I. Spectral estimates.*, Comm. Partial Differential Equations, **19** (1994), No. 9-10, 1397–1447.
- [2] N. Benkirane, *Propriété d'indice en théorie Holderienne pour des opérateurs elliptiques dans  $R^n$* , CRAS, **307**, Série I (1988), 577–580.
- [3] L.A. Caffarelli, N.E. Muler, *An  $L^\infty$  bound for solutions of the Cahn-Hilliard equation*, Arch. Rational Mech. Anal., **133** (1995), No. 2, 129–144
- [4] H.L. Cycon, R.G. Froese, W. Kirsch, B. Simon, *Schrödinger Operators with Application to Quantum Mechanics and Global Geometry*, Springer-Verlag, Berlin (1987).
- [5] A. Ducrot, M. Marion, V. Volpert, *Systemes de réaction-diffusion sans propriété de Fredholm*, CRAS, **340** (2005), 659–664.
- [6] P. Howard, *Spectral analysis of stationary solutions of the Cahn-Hilliard equation*, Adv. Differential Equations, **14** (2009), No. 1-2, 87–120.
- [7] A. Ducrot, M. Marion, V. Volpert, *Reaction-diffusion problems with non Fredholm operators*, Advances Diff. Equations, **13** (2008), No. 11-12, 1151–1192.
- [8] B.L.G. Jonsson, M. Merkli, I.M. Sigal, F. Ting, *Applied Analysis*, In preparation.
- [9] T. Kato, *Wave operators and similarity for some non-selfadjoint operators*, Math. Ann., **162** (1965/1966), 258–279.
- [10] M.D. Korzec, P.L. Evans, A. Münch, B. Wagner, *Stationary solutions of driven fourth- and sixth-order Cahn-Hilliard-type equations*, SIAM J. Appl. Math. , **69** (2008), No. 2, 348–374
- [11] E. Lieb, M. Loss, *Analysis. Graduate Studies in Mathematics*, **14**, American Mathematical Society, Providence (1997).

- [12] M. Reed, B. Simon, *Methods of Modern Mathematical Physics, III: Scattering Theory*, Academic Press (1979).
- [13] I. Rodnianski, W. Schlag, *Time decay for solutions of Schrödinger equations with rough and time-dependent potentials*, *Invent. Math.*, **155** (2004), No. 3, 451–513.
- [14] V. Volpert, B. Kazmierczak, M. Massot, Z. Peradzynski, *Solvability conditions for elliptic problems with non-Fredholm operators*, *Appl. Math.*, **29** (2002), No. 2, 219–238.
- [15] V. Vougalter, V. Volpert, *Solvability conditions for some non Fredholm operators*. To appear in: *Proc. Edinb. Math. Soc.*, <http://hal.archives-ouvertes.fr/hal-00362446/fr/>
- [16] V. Vougalter, V. Volpert. *On the solvability conditions for some non Fredholm operators*, *Int. J. Pure Appl. Math.*, **60** (2010), No. 2, 169–191.
- [17] V. Vougalter, V. Volpert. *On the solvability conditions for the diffusion equation with convection terms*. To appear in: *Commun. Pure Appl. Anal.*
- [18] V. Vougalter, V. Volpert. *Solvability relations for some non Fredholm operators*. To appear in: *Int. Electron. J. Pure Appl. Math.*