

# POINTWISE ESTIMATES AND RIGIDITY RESULTS FOR POSSIBLY SINGULAR AND DEGENERATE ELLIPTIC PDES WITH GENERAL NONLINEARITIES

ALBERTO FARINA<sup>1,2</sup>  
ENRICO VALDINOCI<sup>1,3,4</sup>

ABSTRACT. We prove pointwise gradient bounds for entire solutions of PDEs of the form

$$\mathcal{L}u(x) = \psi(x, u(x), \nabla u(x)),$$

where  $\mathcal{L}$  is an elliptic operator (possibly singular or degenerate).

Thus, we obtain some Liouville type rigidity results. Some classical results of J. Serrin are also recovered as particular cases of our approach.

## 1. INTRODUCTION

A long-established topic in partial differential equations is the study of entire (i.e. reasonably smooth and defined in the whole of  $\mathbb{R}^n$ ) solutions of the equation

$$(1) \quad \mathcal{L}u(x) = \psi(x, u(x), \nabla u(x)) \quad \text{for any } x \in \mathbb{R}^n.$$

In order to obtain regularity and rigidity results, and keeping in mind the important physical applications covered by such models, one considers the case in which  $\mathcal{L}$  is an elliptic operator (possibly with singularities or degeneracies). Typical examples are: the Laplacian operator, in which

$$(2) \quad \mathcal{L}u = \Delta u,$$

the  $p$ -Laplacian, in which

$$(3) \quad \mathcal{L}u = \operatorname{div}(|\nabla u|^{p-2} \nabla u),$$

and the mean curvature operator, namely

$$\mathcal{L}u = \operatorname{div} \left( \frac{\nabla u}{\sqrt{1 + |\nabla u|^2}} \right).$$

Of course, the equation in (3) boils down to the one in (2) when  $p = 2$ . The main feature of all this operators is that they induce some kind of regularity on the solutions, typically in Hölder and Sobolev spaces. On the other hand, a more recent regularity approach focuses on pointwise estimates, to wit, for instance, the gradient of the solution is not only bounded with respect to

---

<sup>1</sup>LAMFA – CNRS UMR 6140 – Université de Picardie Jules Verne – Faculté des Sciences – 33, rue Saint-Leu – 80039 Amiens CEDEX 1, France

<sup>2</sup>Email: [alberto.farina@u-picardie.fr](mailto:alberto.farina@u-picardie.fr)

<sup>3</sup>Università di Roma Tor Vergata – Dipartimento di Matematica – via della ricerca scientifica, 1 – I-00133 Rome, Italy

<sup>4</sup>Email: [enrico@math.utexas.edu](mailto:enrico@math.utexas.edu)

some Hölder or Sobolev norm, but at any point too. These type of pointwise bounds give a very good local control on the solutions and they can be used to obtain further rigidity and symmetry properties.

As far as we know, the idea of dealing with pointwise bounds may be traced back to [Ber27], where it was introduced the idea to look at the equation (or the variational inequality) satisfied by the gradient of the solution, and to deduce universal bounds from that, via the Maximum Principle. This classical approach was used in [Ser72], where new and powerful ideas were introduced in order to prove that entire solutions, under suitable assumptions, need to be constant, thus obtaining important extensions of the so-called Liouville Theorem for harmonic functions.

The original idea of [Ber27] has then been extended and modified in several ways (see, among the others, [Pay76, Spe81, Mod85, CGS94, Far07]) and used for a detailed classification of entire solutions in many cases of interest. In particular, instead of looking at the variational inequality satisfied by the gradient only, it has become relevant to look at a more general variational inequality involving the analogue of the Hamiltonian (or Lagrangian) function in the dynamical system framework. The pointwise estimates obtained in this way are therefore somewhat more precise, since they better take into account the particular features of the nonlinearities involved, and they may be seen as the generalization of the Conservation of Energy Principle to the PDE setting.

The case in which the solution is not entire, but defined on a proper domain of  $\mathbb{R}^n$  has been recently studied in [FV10a, FV10b, CFV09]. In this case, the geometry of the domain (and, in particular, its mean curvature) turn out to play an important role. The situation arising for an elliptic PDE in a compact Riemannian manifold has been considered in [FV11].

The scope of this paper is to deal with entire solutions of equation (1) in a unified framework and under very general assumptions (in particular, we comprise, at the same time, possibly singular and degenerate operators, and gradient dependence in the nonlinearity), obtaining both pointwise gradient estimates and rigidity results, and also recovering some classical results (such as the ones in [Ser72]) as particular cases. For this, we consider the following PDE in divergence form:

$$(4) \quad \operatorname{div}(\Phi'(|\nabla u|^2)\nabla u) = f(u) + g(x, u, \nabla u).$$

Here  $u \in C^1(\mathbb{R}^n)$ ,  $f \in C^1(\mathbb{R})$ ,  $g \in C^1(\mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^n) \cap L^\infty(\mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^n)$ , and the notation  $g := g(x, u, \zeta)$  with  $(x, u, \zeta) \in \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^n$  will be often used. The PDE in (4) is intended in the weak sense and we suppose that  $\Phi$  satisfies the possibly singular or degenerate elliptic conditions listed below.

We recall that PDEs with gradient dependence, of which (4) comprises a quite general setting, are a classical topic of research, see e.g. [DFSU09], and in fact, even ODEs with gradient dependence have been the object of intense study, see [Har02, DFU08] and references therein. Differently from many classical approaches, here we obtain pointwise estimates on the gradient of the solution, and not only estimates in the  $L^\infty$ -norm.

We assume that  $\Phi \in C_{loc}^{3,\alpha}((0, +\infty)) \cap C([0, +\infty))$  for some  $\alpha \in (0, 1)$ , and that  $\Phi(0) = 0$ .

We define

$$(5) \quad a_{ij}(\sigma) := 2\Phi''(|\sigma|^2)\sigma_i\sigma_j + \Phi'(|\sigma|^2)\delta_{ij}.$$

and we suppose that one of the following conditions is satisfied:

**Assumption (A).** There exist  $p > 1$ ,  $a \geq 0$  and  $c_1, c_2 > 0$  such that for any  $\sigma, \xi \in \mathbb{R}^n \setminus \{0\}$

$$(6) \quad c_1(a + |\sigma|)^{p-2} \leq \Phi'(|\sigma|^2) \leq c_2(a + |\sigma|)^{p-2}$$

and

$$(7) \quad c_1(a + |\sigma|)^{p-2} |\xi|^2 \leq \sum_{i,j=1}^n a_{ij}(\sigma) \xi_i \xi_j \leq c_2(a + |\sigma|)^{p-2} |\xi|^2.$$

**Assumption (B).**  $\Phi \in C^1([0, +\infty))$ , and there exist  $c_1, c_2 > 0$  such that for any  $\sigma \in \mathbb{R}^n$

$$(8) \quad c_1(1 + |\sigma|)^{-1} \leq \Phi'(|\sigma|^2) \leq c_2(1 + |\sigma|)^{-1}$$

and

$$(9) \quad c_1(1 + |\sigma|)^{-1} |\xi'|^2 \leq \sum_{i,j=1}^n a_{ij}(\sigma) \xi_i \xi_j \leq c_2(1 + |\sigma|)^{-1} |\xi'|^2,$$

for any  $\xi' = (\xi, \xi_{n+1}) \in \mathbb{R}^{n+1}$  which is orthogonal to  $(-\sigma, 1) \in \mathbb{R}^{n+1}$ .

The above Assumptions (A) and (B) are classical: they agree, for instance, with the ones of [CGS94], and examples of functional satisfying the above conditions are the  $p$ -Laplacian (with  $p \in (1, +\infty)$ ) and the mean curvature operators – which correspond to the cases

$$\Phi(r) := \frac{2}{p} r^{p/2} \quad \text{and} \quad \Phi(r) := 2\sqrt{1+r} - 2,$$

respectively.

The simplest case of our result can be stated in the following way:

**Theorem 1.1.** *Let  $u \in C^1(\mathbb{R}^n) \cap W^{1,\infty}(\mathbb{R}^n)$  be a weak solution in the whole of  $\mathbb{R}^n$  of one of the following equations: either*

$$(10) \quad \begin{cases} \operatorname{div}(\Phi'(|\nabla u|^2) \nabla u) = h(u) + c(x) \cdot \nabla u \\ \text{with } h \in C^1(\mathbb{R}), c \in C^1(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n), \\ h' \geq 0 \text{ and } \left\{ \frac{\partial c_i}{\partial x_j} \right\}_{\{i,j=1,\dots,n\}} \text{ a nonnegative matrix,} \end{cases}$$

or

$$(11) \quad \begin{cases} \operatorname{div}(\Phi'(|\nabla u|^2) \nabla u) = f(u) + g(\nabla u) \\ \text{with } f \in C^1(\mathbb{R}), g \in C^1(\mathbb{R}^n) \text{ and } fg \leq 0. \end{cases}$$

Let  $F := 0$  if (10) holds, and  $F$  be a primitive<sup>1</sup> of  $f$  with  $F \geq 0$  if (11) holds. Then,

$$(12) \quad 2\Phi'(|\nabla u(x)|^2) |\nabla u(x)|^2 - \Phi(|\nabla u(x)|^2) \leq 2F(u(x)) \quad \text{for any } x \in \mathbb{R}^n.$$

More generally, we prove the following results:

**Theorem 1.2.** *Let  $u \in C^1(\mathbb{R}^n) \cap W^{1,\infty}(\mathbb{R}^n)$  be a weak solution of (4) in the whole of  $\mathbb{R}^n$ , with  $f \in C^1(\mathbb{R})$ ,  $g \in C^1(\mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^n) \cap L^\infty(\mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^n)$ .*

*Let  $F$  be a primitive of  $f$ , with  $F \geq 0$ . Let*

$$(13) \quad \mathcal{R}(x) := -\frac{2f(u)g(x, u, \nabla u)|\nabla u|^2}{\Phi'(|\nabla u|^2)} + 2|\nabla u|^2 \nabla_x g(x, u, \nabla u) \cdot \nabla u + 2|\nabla u|^4 g_u(x, u, \nabla u)$$

---

<sup>1</sup>We observe that, since  $u$  is bounded, we can find a primitive of  $f$  that is non-negative in the range of  $u$ .

and assume that

$$(14) \quad \mathcal{R}(x) \geq 0 \text{ for any } x \in \mathbb{R}^n.$$

Then,

$$(15) \quad 2\Phi'(|\nabla u(x)|^2)|\nabla u(x)|^2 - \Phi(|\nabla u(x)|^2) \leq 2F(u(x)) \quad \text{for any } x \in \mathbb{R}^n.$$

**Theorem 1.3.** *Let the assumptions of Theorem 1.2 hold.*

*Also, if Assumption (A) holds with  $p \geq 2$ , assume that for any  $\mu \in \{F = 0\}$  we have  $F(r) = O(|r - \mu|^p)$ .*

*Then, if there exists  $x_o \in \mathbb{R}^n$  for which  $F(u(x_o)) = 0$ , we have that  $u(x) = u(x_o)$  for any  $x \in \mathbb{R}^n$ .*

Theorem 1.2 is a pointwise estimate on the gradient of the solution and Theorem 1.3 is a rigidity result of Liouville type. When  $g := 0$ , similar results have been obtained by [Mod85] in the semilinear case  $\Phi(r) := r$  and by [CGS94] under Assumptions (A) or (B). Then, Theorems 1.2 and 1.3 here may be seen as extensions of the works of [Mod85, CGS94] to the case of more complicated nonlinearities involving  $g(x, u, \nabla u)$ .

We remark that condition (14), although artificial at first glance, comprises many cases of interest, such as:

- $f := 0$ ,  $g(x, u, \zeta) := h(u) + c(x) \cdot \zeta$ , with  $h' \geq 0$  and  $\{\frac{\partial c_i}{\partial x_j}\}_{\{i,j=1,\dots,n\}}$  a nonnegative matrix;
- $g = g(\zeta)$  and  $fg \leq 0$ .

These cases, which are the ones presented in Theorem 1.1, may also be seen as extensions of some of the results of [Ser72] to the case in which the operator is not uniformly elliptic and in divergence form, and techniques we use are different (see also Appendix A).

We also remark that, in its generality, Theorem 1.2 is new, to the best of our knowledge, even in the semilinear case  $\Phi(r) := r$ . See also [FV10b, FV11, CFV09] for related works on proper domains and on manifolds.

**Example 1.1.** If condition (14) is dropped, then (15) may not hold, as the following example shows. Let  $n := 1$ ,  $u(x) := \arctan(x)$ ,  $\Phi(r) := r$ ,  $f := 0$ ,  $F := 0$  and  $g(x, u, \zeta) := -2x/(1+x^2)^2$ . Then we see that  $u'' = f + g$ , hence (4) is satisfied, that  $\mathcal{R} = 2|u'(x)|^2 g'(x)u'(x) = 4(3x^2 - 1)/(1+x^2)^3$ , hence (14) does not hold, and (15) is violated because  $u$  is not constant.

Example 1.1 also shows that (15) is quite a strong estimate, since the gradient of the solution is estimated (at all points, and not only in the average) by something that depends only on  $f$ , not on  $g$ .

The proof of Theorems 1.2 and 1.3 will be based on a long and delicate computation, detailed in Section 2, and on the proofs of the results of [CGS94], as discussed in Section 3.

## 2. P-FUNCTION COMPUTATIONS

Let

$$(16) \quad \Lambda(s) := 2s\Phi''(s) + \Phi'(s),$$

and

$$(17) \quad d_{ij}(\sigma) := \frac{a_{ij}(\sigma)}{\Lambda(|\sigma|^2)}.$$

We define

$$(18) \quad P(u, x) = 2\Phi'(|\nabla u(x)|^2)|\nabla u(x)|^2 - \Phi(|\nabla u(x)|^2) - 2F(u(x)).$$

The main idea driving the following computation comes from some classical works such as [Pay76, Spe81, CGS94] in which it is shown that some suitable  $P$ -function solves an elliptic PDE (at least at nonsingular points). The Maximum Principle then provides an estimate on  $P$ , which, in turn, would give the desired result.

With this goal in mind, we pursue the following result (which, for  $g := 0$  reduces to formula (2.7) of [CGS94]):

**Lemma 2.1.** *Let  $\Omega$  be an open subset of  $\mathbb{R}^n$ . Let  $u \in C^1(\Omega)$  be a solution of (4) in  $\Omega$ , with  $\nabla u \neq 0$  in  $\Omega$ .*

Let

$$B_i(x) := -2 \frac{f(u)}{\Lambda(|\nabla u|^2)} \left( 1 + \frac{|\nabla u|^2 \Phi''(|\nabla u|^2)}{\Phi'(|\nabla u|^2)} \right) \frac{\partial u}{\partial x_i} - \frac{|\nabla u|^2}{\Lambda(|\nabla u|^2)} \frac{\partial g}{\partial p_i}(x, u, \nabla u).$$

Then,

$$(19) \quad \sum_{ij} |\nabla u|^2 \frac{\partial}{\partial x_j} \left( d_{ij}(\nabla u) \frac{\partial P}{\partial x_i} \right) + \sum_i B_i \frac{\partial P}{\partial x_i} \geq \frac{|\nabla P|^2}{2\Lambda(|\nabla u|^2)} + \mathcal{R} \quad \text{weakly in } \Omega.$$

*Proof.* First, we remark that, by our assumptions, the map  $r \mapsto 2\Phi'(r)r - \Phi(r)$  is invertible. We call  $\Psi$  its inverse. Notice that

$$(20) \quad \Psi(P(u, x) + 2F(u(x))) = |\nabla u(x)|^2.$$

Moreover, by the definition of  $\Psi$  and (16),

$$1 = \frac{d}{dr} \left( \Psi \left( 2\Phi'(r)r - \Phi(r) \right) \right) = \Psi' \left( 2\Phi'(r)r - \Phi(r) \right) \Lambda(r),$$

hence

$$(21) \quad \Psi' \left( 2\Phi'(|\nabla u|^2)|\nabla u|^2 - \Phi(|\nabla u|^2) \right) = \frac{1}{\Lambda(|\nabla u|^2)}.$$

Now, differentiating (18) and recalling (16), we see that

$$(22) \quad \frac{\partial P}{\partial x_i} = 2\Lambda(|\nabla u|^2) \sum_k \frac{\partial^2 u}{\partial x_i \partial x_k} \frac{\partial u}{\partial x_k} - 2f(u) \frac{\partial u}{\partial x_i}$$

hence, recalling (17),

$$(23) \quad \begin{aligned} \sum_{ij} \frac{\partial}{\partial x_j} \left( d_{ij}(\nabla u) \frac{\partial P}{\partial x_i} \right) &= -2 \sum_{ij} \frac{\partial}{\partial x_j} \left( f(u) d_{ij}(\nabla u) \frac{\partial u}{\partial x_i} \right) + \\ &+ 2 \sum_{ijk} \frac{\partial}{\partial x_j} \left( a_{ij}(\nabla u) \frac{\partial^2 u}{\partial x_i \partial x_k} \right) \frac{\partial u}{\partial x_k} + \\ &+ 2 \sum_{ijk} a_{ij}(\nabla u) \frac{\partial^2 u}{\partial x_i \partial x_k} \frac{\partial^2 u}{\partial x_j \partial x_k}. \end{aligned}$$

Also, (5) gives that

$$(24) \quad \frac{\partial a_{ij}}{\partial \sigma_\ell}(\sigma) = \frac{\partial a_{\ell j}}{\partial \sigma_i}(\sigma)$$

and, by (4),

$$(25) \quad \sum_{ij} a_{ij}(\nabla u) \frac{\partial^2 u}{\partial x_i \partial x_j} = f(u) + g(x, u, \nabla u).$$

Therefore, by (24) and (25), for any fixed  $k$ ,

$$(26) \quad \begin{aligned} \sum_{ijk} \frac{\partial}{\partial x_j} \left( a_{ij}(\nabla u) \frac{\partial^2 u}{\partial x_i \partial x_k} \right) &= \sum_{ijk} \frac{\partial}{\partial x_k} \left( a_{ij}(\nabla u) \frac{\partial^2 u}{\partial x_i \partial x_j} \right) = \\ &= f'(u) \frac{\partial u}{\partial x_k} + g_{x_k} + g_u \frac{\partial u}{\partial x_k} + \sum_i g_{p_i} \frac{\partial^2 u}{\partial x_i \partial x_k}, \end{aligned}$$

where the notation  $g = g(x, u, \zeta)$  has been used.

From (23) and (26), we gather

$$(27) \quad \begin{aligned} \sum_{ij} \frac{\partial}{\partial x_j} \left( d_{ij}(\nabla u) \frac{\partial P}{\partial x_i} \right) &= \\ &= 2f'(u) \sum_k \frac{\partial u}{\partial x_k} \frac{\partial u}{\partial x_k} + 2 \sum_{ijk} a_{ij}(\nabla u) \frac{\partial^2 u}{\partial x_i \partial x_k} \frac{\partial^2 u}{\partial x_j \partial x_k} - \\ &- 2 \sum_{ij} f'(u) d_{ij}(\nabla u) \frac{\partial u}{\partial x_i} \frac{\partial u}{\partial x_j} - 2f(u) \sum_{ij} \frac{\partial}{\partial x_j} \left( d_{ij}(\nabla u) \frac{\partial u}{\partial x_i} \right) + \\ &+ 2 \sum_k \left[ g_{x_k} + g_u \frac{\partial u}{\partial x_k} + \sum_i g_{p_i} \frac{\partial^2 u}{\partial x_i \partial x_k} \right] \frac{\partial u}{\partial x_k}. \end{aligned}$$

Furthermore, from (16) and (17), we obtain

$$\begin{aligned} f'(u) \sum_k \frac{\partial u}{\partial x_k} \frac{\partial u}{\partial x_k} - \sum_{ij} f'(u) d_{ij}(\nabla u) \frac{\partial u}{\partial x_i} \frac{\partial u}{\partial x_j} &= \\ &= f'(u) \left[ |\nabla u|^2 - \frac{\Phi'(|\nabla u|^2) |\nabla u|^2 + 2\Phi''(|\nabla u|^2) |\nabla u|^4}{\Lambda(|\nabla u|^2)} \right] = \\ &= f'(u) [|\nabla u|^2 - |\nabla u|^2] = 0. \end{aligned}$$

Plugging this into (27), we conclude that

$$\begin{aligned}
(28) \quad & \sum_{ij} \frac{\partial}{\partial x_j} \left( d_{ij}(\nabla u) \frac{\partial P}{\partial x_i} \right) = \\
& = 2 \sum_{ijk} a_{ij}(\nabla u) \frac{\partial^2 u}{\partial x_i \partial x_k} \frac{\partial^2 u}{\partial x_j \partial x_k} - 2f(u) \sum_{ij} \frac{\partial}{\partial x_j} \left( d_{ij}(\nabla u) \frac{\partial u}{\partial x_i} \right) + \\
& + 2 \sum_k \left[ g_{x_k} + g_u \frac{\partial u}{\partial x_k} + \sum_i g_{p_i} \frac{\partial^2 u}{\partial x_i \partial x_k} \right] \frac{\partial u}{\partial x_k}.
\end{aligned}$$

Also, from (17) and (25),

$$\sum_{ij} d_{ij}(\nabla u) \frac{\partial^2 u}{\partial x_i \partial x_j} = \frac{f(u) + g(x, u, \nabla u)}{\Lambda(|\nabla u|^2)},$$

and so (28) becomes

$$\begin{aligned}
(29) \quad & \sum_{ij} \frac{\partial}{\partial x_j} \left( d_{ij}(\nabla u) \frac{\partial P}{\partial x_i} \right) = \\
& = 2 \sum_{ijk} a_{ij}(\nabla u) \frac{\partial^2 u}{\partial x_i \partial x_k} \frac{\partial^2 u}{\partial x_j \partial x_k} - \\
& - 2f(u) \sum_{ij} \frac{\partial}{\partial x_j} d_{ij}(\nabla u) \frac{\partial u}{\partial x_i} - 2 \frac{f(u) [f(u) + g(x, u, \nabla u)]}{\Lambda(|\nabla u|^2)} + \\
& + 2 \sum_k \left[ g_{x_k} + g_u \frac{\partial u}{\partial x_k} + \sum_i g_{p_i} \frac{\partial^2 u}{\partial x_i \partial x_k} \right] \frac{\partial u}{\partial x_k}.
\end{aligned}$$

Moreover, making use of (5), (16) and (17), we obtain

$$\begin{aligned}
(30) \quad & \sum_{ij} \frac{\partial}{\partial x_j} d_{ij}(\nabla u) \frac{\partial u}{\partial x_i} = \\
& = \sum_{ij} \frac{\partial u}{\partial x_i} \frac{\partial}{\partial x_j} \frac{2\Phi''(|\nabla u|^2) \frac{\partial u}{\partial x_i} \frac{\partial u}{\partial x_j} + \Phi'(|\nabla u|^2) \delta_{ij}}{2|\nabla u|^2 \Phi''(|\nabla u|^2) + \Phi'(|\nabla u|^2)} = \\
& = \sum_{ij} \frac{\partial u}{\partial x_i} \frac{\left[ 4\Phi''' \sum_k \frac{\partial u}{\partial x_k} \frac{\partial^2 u}{\partial x_j \partial x_k} \frac{\partial u}{\partial x_i} \frac{\partial u}{\partial x_j} + 2\Phi'' \frac{\partial^2 u}{\partial x_j \partial x_i} \frac{\partial u}{\partial x_j} + 2\Phi'' \frac{\partial u}{\partial x_i} \frac{\partial^2 u}{\partial x_j^2} + 2\Phi'' \sum_k \frac{\partial u}{\partial x_k} \frac{\partial^2 u}{\partial x_j \partial x_k} \delta_{ij} \right]}{2|\nabla u|^2 \Phi'' + \Phi'} \\
& - \sum_{ij} \frac{\partial u}{\partial x_i} \frac{(2\Phi'' \frac{\partial u}{\partial x_i} \frac{\partial u}{\partial x_j} + \Phi' \delta_{ij}) \left[ 6\Phi'' + 4|\nabla u|^2 \Phi''' \right] \sum_k \frac{\partial u}{\partial x_k} \frac{\partial^2 u}{\partial x_j \partial x_k}}{(2|\nabla u|^2 \Phi'' + \Phi')^2} = \\
& = 2 \frac{\Phi''(|\nabla u|^2)}{\Lambda(|\nabla u|^2)} \left( |\nabla u|^2 \Delta u - \sum_{ij} \frac{\partial^2 u}{\partial x_i \partial x_j} \frac{\partial u}{\partial x_i} \frac{\partial u}{\partial x_j} \right).
\end{aligned}$$

Also, from (5) and (25),

$$\begin{aligned}
& f(u) + g(x, u, \nabla u) = \\
& = \sum_{ij} \left( 2\Phi''(|\nabla u|^2) \frac{\partial u}{\partial x_i} \frac{\partial u}{\partial x_j} + \Phi'(|\nabla u|^2) \delta_{ij} \right) \frac{\partial^2 u}{\partial x_i \partial x_j} = \\
& = 2\Phi''(|\nabla u|^2) \sum_{ij} \frac{\partial^2 u}{\partial x_i \partial x_j} \frac{\partial u}{\partial x_i} \frac{\partial u}{\partial x_j} + \Phi'(|\nabla u|^2) \Delta u,
\end{aligned}$$

from which we obtain

$$\Delta u = \frac{f(u) + g(x, u, \nabla u)}{\Phi'(|\nabla u|^2)} - 2 \frac{\Phi''(|\nabla u|^2)}{\Phi'(|\nabla u|^2)} \sum_{ij} \frac{\partial^2 u}{\partial x_i \partial x_j} \frac{\partial u}{\partial x_i} \frac{\partial u}{\partial x_j}.$$

Therefore, recalling also (22), we write (30) as

$$\begin{aligned}
(31) \quad & \sum_{ij} \frac{\partial}{\partial x_j} d_{ij}(\nabla u) \frac{\partial u}{\partial x_j} = \\
& = 2 \frac{\Phi''(|\nabla u|^2)}{\Phi'(|\nabla u|^2) \Lambda(|\nabla u|^2)} \left[ (f + g) |\nabla u|^2 - \Lambda(|\nabla u|^2) \sum_{ij} \frac{\partial^2 u}{\partial x_i \partial x_j} \frac{\partial u}{\partial x_i} \frac{\partial u}{\partial x_j} \right] = \\
& = - \frac{\Phi''(|\nabla u|^2)}{\Phi'(|\nabla u|^2) \Lambda(|\nabla u|^2)} \sum_i \frac{\partial P}{\partial x_i} \frac{\partial u}{\partial x_i} + \\
& + \frac{2g\Phi''(|\nabla u|^2) |\nabla u|^2}{\Phi'(|\nabla u|^2) \Lambda(|\nabla u|^2)}.
\end{aligned}$$



Thus, exploiting (29),

$$\begin{aligned}
(32) \quad & \sum_{ij} \frac{\partial}{\partial x_j} \left( d_{ij}(\nabla u) \frac{\partial P}{\partial x_i} \right) \\
&= 2f \frac{\Phi''(|\nabla u|^2)}{\Phi'(|\nabla u|^2)\Lambda(|\nabla u|^2)} \sum_i \frac{\partial P}{\partial x_i} \frac{\partial u}{\partial x_i} + \\
&+ 2 \sum_{ijk} a_{ij}(\nabla u) \frac{\partial^2 u}{\partial x_i \partial x_k} \frac{\partial^2 u}{\partial x_j \partial x_k} - 2 \frac{f[f+g]}{\Lambda(|\nabla u|^2)} + \\
&+ 2 \sum_k \left[ g_{x_k} + g_u \frac{\partial u}{\partial x_k} + \sum_i g_{p_i} \frac{\partial^2 u}{\partial x_i \partial x_k} \right] \frac{\partial u}{\partial x_k} - \\
&- \frac{4fg\Phi''(|\nabla u|^2)|\nabla u|^2}{\Phi'(|\nabla u|^2)\Lambda(|\nabla u|^2)}.
\end{aligned}$$

Now we set

$$z_k = \sum_i \frac{\partial^2 u}{\partial x_i \partial x_k} \frac{\partial u}{\partial x_i},$$

and we use Schwarz Inequality to see that

$$|z_k| \leq \sqrt{\sum_i \left( \frac{\partial^2 u}{\partial x_i \partial x_k} \right)^2} \sqrt{\sum_i \left( \frac{\partial u}{\partial x_i} \right)^2}$$

and so

$$\begin{aligned}
& \sum_{ijk} \frac{\partial^2 u}{\partial x_i \partial x_k} \frac{\partial u}{\partial x_i} \frac{\partial^2 u}{\partial x_j \partial x_k} \frac{\partial u}{\partial x_j} = \sum_k z_k^2 \leq \\
& \leq \sum_k \left( \sum_i \left( \frac{\partial^2 u}{\partial x_i \partial x_k} \right)^2 \right) \left( \sum_i \left( \frac{\partial u}{\partial x_i} \right)^2 \right) = \\
& = |\nabla u|^2 \sum_{ik} \left( \frac{\partial^2 u}{\partial x_i \partial x_k} \right)^2.
\end{aligned}$$

This and (5) give that

$$\begin{aligned}
(33) \quad & \sum_{ijk} a_{ij}(\nabla u) \frac{\partial^2 u}{\partial x_i \partial x_k} \frac{\partial^2 u}{\partial x_j \partial x_k} \geq \\
& \geq \sum_{ijk} \frac{\Phi'}{|\nabla u|^2} \frac{\partial^2 u}{\partial x_i \partial x_k} \frac{\partial u}{\partial x_i} \frac{\partial^2 u}{\partial x_j \partial x_k} \frac{\partial u}{\partial x_j} + \\
& + 2\Phi'' \sum_{ijk} \frac{\partial^2 u}{\partial x_i \partial x_k} \frac{\partial u}{\partial x_i} \frac{\partial^2 u}{\partial x_j \partial x_k} \frac{\partial u}{\partial x_j} = \\
& = \frac{\Lambda}{|\nabla u|^2} \sum_{ijk} \frac{\partial^2 u}{\partial x_i \partial x_k} \frac{\partial u}{\partial x_i} \frac{\partial^2 u}{\partial x_j \partial x_k} \frac{\partial u}{\partial x_j}.
\end{aligned}$$

Moreover, by (22),

$$\sum_{ijk} \frac{\partial^2 u}{\partial x_i \partial x_k} \frac{\partial u}{\partial x_i} \frac{\partial^2 u}{\partial x_j \partial x_k} \frac{\partial u}{\partial x_j} = \frac{1}{4\Lambda^2} \sum_k \left( \frac{\partial P}{\partial x_k} + 2f \frac{\partial u}{\partial x_k} \right)^2$$

and so (33) becomes

$$\begin{aligned} \sum_{ijk} a_{ij}(\nabla u) \frac{\partial^2 u}{\partial x_i \partial x_k} \frac{\partial^2 u}{\partial x_j \partial x_k} &\geq \frac{1}{4|\nabla u|^2 \Lambda} \sum_k \left( \frac{\partial P}{\partial x_k} + 2f \frac{\partial u}{\partial x_k} \right)^2 = \\ &= \frac{|\nabla P|^2}{4|\nabla u|^2 \Lambda} + \frac{f \sum_i \frac{\partial u}{\partial x_i} \frac{\partial P}{\partial x_i}}{|\nabla u|^2 \Lambda} + \frac{f^2}{\Lambda}. \end{aligned}$$

By substituting this in (32), we obtain

$$\begin{aligned} (34) \quad \sum_{ij} \frac{\partial}{\partial x_j} \left( d_{ij}(\nabla u) \frac{\partial P}{\partial x_i} \right) - \frac{2f}{|\nabla u|^2 \Lambda} \left( 1 + \frac{|\nabla u|^2 \Phi''}{\Phi'} \right) \sum_i \frac{\partial P}{\partial x_i} \frac{\partial u}{\partial x_i} &\geq \\ &\geq \frac{|\nabla P|^2}{2|\nabla u|^2 \Lambda} - \frac{2fg}{\Phi'} + 2 \sum_k \left[ g_{x_k} + g_u \frac{\partial u}{\partial x_k} + \sum_i g_{p_i} \frac{\partial^2 u}{\partial x_i \partial x_k} \right] \frac{\partial u}{\partial x_k}. \end{aligned}$$

Now, we use (20), according to which, for  $i$  fixed,

$$2 \sum_k \frac{\partial^2 u}{\partial x_i \partial x_k} \frac{\partial u}{\partial x_k} = \Psi'(P + 2F) \left( \frac{\partial P}{\partial x_i} + 2f \frac{\partial u}{\partial x_i} \right).$$

Accordingly,

$$2 \sum_{ki} g_{p_i} \frac{\partial^2 u}{\partial x_i \partial x_k} \frac{\partial u}{\partial x_k} = \Psi'(P + 2F) \sum_i g_{p_i} \left( \frac{\partial P}{\partial x_i} + 2f \frac{\partial u}{\partial x_i} \right).$$

Plugging this in (34) and recalling (21) we obtain the desired result.  $\square$

### 3. PROOF OF THEOREMS 1.2 AND 1.3

We observe that, since  $u \in C^1(\mathbb{R}^n) \cap W^{1,\infty}(\mathbb{R}^n)$ , our assumptions on  $f$  and  $g$  imply that  $u \in C_{\text{loc}}^{1,\alpha}(\mathbb{R}^n)$  and the family of all the translations of  $u$  is relatively compact in  $C_{\text{loc}}^{1,\alpha}(\mathbb{R}^n)$ , for a suitable  $\alpha \in (0, 1)$ , both under assumptions (A) and (B) (see, for instance, [DiB83, Tol84, GT01]). Since  $\mathcal{R} \geq 0$ , we know by Lemma 2.1 that

$$\sum_{ij} \frac{\partial}{\partial x_j} \left( d_{ij}(\nabla u) \frac{\partial P}{\partial x_i} \right) + \frac{\nabla B \cdot \nabla P}{|\nabla u|^2} \geq 0$$

weakly in  $\{\nabla u \neq 0\}$ .

Then, we can repeat the arguments in the proofs of Theorems 1.6 and 1.8 of [CGS94] (see pages 1464–1466 there) and obtain our Theorems 1.2 and 1.3.  $\square$

APPENDIX A. RECOVERING SOME IMPORTANT RESULTS OF [Ser72] AS PARTICULAR CASES OF OUR RESULTS

The purpose of this appendix is to recover some important results of [Ser72]. More precisely, from (1) and (2) in [Ser72], one has the following results:

**Theorem A.1.** [Page 348 in [Ser72]] Let  $g \in C^1(\mathbb{R} \times \mathbb{R}^n)$ . Let  $u \in C^3(\mathbb{R}^n) \cap W^{1,\infty}(\mathbb{R}^n)$  be a solution of

$$(35) \quad \Delta u = g(u, \nabla u)$$

in  $\mathbb{R}^n$ , with

$$(36) \quad g_u \geq 0.$$

Then  $u$  is constant.

**Theorem A.2.** [Page 349 in [Ser72]] Let  $\psi \in C^1(\mathbb{R})$ . Let  $u \in C^3(\mathbb{R}^n) \cap W^{1,\infty}(\mathbb{R}^n)$  be a solution of

$$(37) \quad (1 + |\nabla u|^2)\Delta u - u_i u_j u_{ij} = \psi(\nabla u)$$

in  $\mathbb{R}^n$ . Then  $u$  is constant.

We can obtain Theorems A.1 and A.2 directly from our Theorems 1.2 and 1.1, respectively (Theorem 1.3 could have also been used) via the following argument. For Theorem A.1, we take  $f := 0$ ,  $F := 0$  and  $\Phi(r) := r$ , hence the left hand side of (15) equals to  $|\nabla u|^2$ . In this case (4) agrees with (35) and  $g$  is independent of  $x$ . Also, by (13) and (36),  $\mathcal{R} = 2|\nabla u|^4 g_u \geq 0$ , so (14) holds true. Thus, since  $F$  vanishes identically, by (15), we have that  $\nabla u$  vanishes identically too, giving a new proof of Theorem A.1.

As for Theorem A.2, we take  $f := 0$ ,  $F := 0$ ,  $g(p) := \psi(p)/(1 + |p|^2)^{3/2}$  and  $\Phi(r) := 2\sqrt{1+r} - 2$ . Then (37) agrees with (11), and the left hand side of (12) is

$$\frac{2\left(\sqrt{1 + |\nabla u|^2} - 1\right)}{\sqrt{1 + |\nabla u|^2}},$$

which is nonnegative, and it vanishes if and only if  $\nabla u = 0$ . Accordingly, since  $F$  vanishes identically, (12) gives that  $\nabla u$  vanishes identically as well, giving a new proof of Theorem A.2.  $\square$

#### ACKNOWLEDGMENTS

This work has been supported by *FIRB* project ‘‘Analysis and Beyond’’ and *GNAMPA* project ‘‘Equazioni nonlineari su variet : propriet  qualitative e classificazione delle soluzioni’’.

#### REFERENCES

- [Ber27] Serge Bernstein.  ber ein geometrisches Theorem und seine Anwendung auf die partiellen Differentialgleichungen vom elliptischen Typus. *Math. Z.*, 26(1):551–558, 1927.
- [CFV09] Diego Castellaneta, Alberto Farina, and Enrico Valdinoci. A pointwise gradient estimate for solutions of singular and degenerate PDEs in possibly unbounded domains with nonnegative mean curvature. *Preprint*, 2009.
- [CGS94] Luis Caffarelli, Nicola Garofalo, and Fausto Segala. A gradient bound for entire solutions of quasi-linear equations and its consequences. *Comm. Pure Appl. Math.*, 47(11):1457–1473, 1994.

- [DFSU09] Djairo G. De Figueiredo, Justino Sánchez, and Pedro Ubilla. Quasilinear equations with dependence on the gradient. *Nonlinear Anal.*, 71(10):4862–4868, 2009.
- [DFU08] Djairo G. De Figueiredo and Pedro Ubilla. Superlinear systems of second-order ODE's. *Nonlinear Anal.*, 68(6):1765–1773, 2008.
- [DiB83] E. DiBenedetto.  $C^{1+\alpha}$  local regularity of weak solutions of degenerate elliptic equations. *Nonlinear Anal.*, 7(8):827–850, 1983.
- [Far07] Alberto Farina. Liouville-type theorems for elliptic problems. In *Handbook of differential equations: stationary partial differential equations. Vol. IV*, Handb. Differ. Equ., pages 61–116. Elsevier/North-Holland, Amsterdam, 2007.
- [FV10a] Alberto Farina and Enrico Valdinoci. Flattening results for elliptic PDEs in unbounded domains with applications to overdetermined problems. *Arch. Ration. Mech. Anal.*, 195(3):1025–1058, 2010.
- [FV10b] Alberto Farina and Enrico Valdinoci. A pointwise gradient estimate in possibly unbounded domains with nonnegative mean curvature. *Adv. Math.*, 225(5):2808–2827, 2010.
- [FV11] Alberto Farina and Enrico Valdinoci. A pointwise gradient bound for elliptic equations on compact manifolds with nonnegative Ricci curvature. *Discrete Contin. Dyn. Syst.*, 30(4):1139–1144, 2011.
- [GT01] David Gilbarg and Neil S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition.
- [Har02] Philip Hartman. *Ordinary differential equations*, volume 38 of *Classics in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2002. Corrected reprint of the second (1982) edition [Birkhäuser, Boston, MA; MR0658490 (83e:34002)], With a foreword by Peter Bates.
- [Mod85] Luciano Modica. A gradient bound and a Liouville theorem for nonlinear Poisson equations. *Comm. Pure Appl. Math.*, 38(5):679–684, 1985.
- [Pay76] L. E. Payne. Some remarks on maximum principles. *J. Analyse Math.*, 30:421–433, 1976.
- [Ser72] James Serrin. Entire solutions of nonlinear Poisson equations. *Proc. London. Math. Soc. (3)*, 24:348–366, 1972.
- [Spe81] René P. Sperb. *Maximum principles and their applications*, volume 157 of *Mathematics in Science and Engineering*. Academic Press Inc. [Harcourt Brace Jovanovich Publishers], New York, 1981.
- [Tol84] Peter Tolksdorf. Regularity for a more general class of quasilinear elliptic equations. *J. Differential Equations*, 51(1):126–150, 1984.