Erratum
Exact Ground State Energy of the Strong-Coupling Polaron

Commun. Math. Phys. **183**, 511-519, (1997)

Elliott H. Lieb[1] and Lawrence E. Thomas[2]

[1]*Departments of Physics and Mathematics, Jadwin Hall, Princeton University, P. O. Box 708, Princeton, New Jersey 08544*

[2]*Department of Mathematics, University of Virginia, Charlottesville, Virginia 22903*

April 25, 1997

We are grateful to Professor Andrey V. Soldatov of the Moscow Steklov Mathematical Institute for calling our attention to an error in our paper. The commutator inequality (8) in our step **I**, namely $|k_j||\langle a_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{x}}\rangle| \leq 2\langle p_j^2\rangle^{1/2}\langle a_{\mathbf{k}}^* a_{\mathbf{k}}\rangle^{1/2}$, is not correct. Rather, the right side of this inequality should be $\langle p_j^2\rangle^{1/2}(\langle a_{\mathbf{k}}^* a_{\mathbf{k}}\rangle^{1/2} + \langle a_{\mathbf{k}} a_{\mathbf{k}}^*\rangle^{1/2})$ or a related expression. The extra factor $\langle a_{\mathbf{k}} a_{\mathbf{k}}^*\rangle^{1/2}$ with the $a_{\mathbf{k}}$ and $a_{\mathbf{k}}^*$ not in normal order generates uncontrolled mischief with, for example, the right side of the ultraviolet bound (10) containing an additional term $\sum_{|\mathbf{k}|\geq K} 1/2 = \infty$.

The situation is easily remedied with the help of the method introduced by Lieb and Yamazaki [14] to obtain the previous rigorous lower bound on the polaron energy. Our main result, (31), is still valid. Indeed, it is improved slightly!

Define the (vector) operator $\mathbf{Z} = (Z_1, Z_2, Z_3)$ with components

$$Z_j = (\frac{4\pi\alpha}{V})^{1/2} \sum_{|\mathbf{k}|\geq K} k_j \frac{a_{\mathbf{k}}}{|\mathbf{k}|^3} e^{i\mathbf{k}\cdot\mathbf{x}}, \quad j = 1, 2, 3. \tag{1}$$

Then the commutator estimate (8) is replaced by

$$-(\frac{4\pi\alpha}{V})^{1/2} \sum_{|\mathbf{k}|\geq K} \left[\langle \frac{a_{\mathbf{k}}}{|\mathbf{k}|} e^{i\mathbf{k}\cdot\mathbf{x}}\rangle + c.c.\right] \equiv -\sum_j \langle [p_j, Z_j - Z_j^*]\rangle$$

$$\leq 2\langle \mathbf{p}^2\rangle^{1/2}\langle -(\mathbf{Z}-\mathbf{Z}^*)^2\rangle^{1/2} \;\leq\; 2\langle \mathbf{p}^2\rangle^{1/2}\langle 2(\mathbf{Z}^*\mathbf{Z}+\mathbf{Z}\mathbf{Z}^*)\rangle^{1/2}$$
$$\leq \varepsilon\langle \mathbf{p}^2\rangle + \frac{2}{\varepsilon}\langle \mathbf{Z}^*\mathbf{Z}+\mathbf{Z}\mathbf{Z}^*\rangle. \qquad (2)$$

Now, each component Z_j can be thought of as a single (unnormalized) oscillator mode having commutator with its adjoint, $[Z_j, Z_j^*] = (4\pi\alpha/V)\sum_{|\mathbf{k}|\geq K} k_j^2|\mathbf{k}|^{-6} \to 2\alpha/3\pi K$; moreover, Z_i and Z_j^* commute for $i \neq j$ (i.e., these modes are orthogonal). Using these facts, we have that

$$\frac{2}{\varepsilon}\langle \mathbf{Z}^*\mathbf{Z}+\mathbf{Z}\mathbf{Z}^*\rangle \;=\; \frac{4}{\varepsilon}\langle \mathbf{Z}^*\mathbf{Z}\rangle + \frac{2}{\varepsilon}\left(\frac{2\alpha}{\pi K}\right)$$
$$\leq \sum_{|\mathbf{k}|\geq K}\langle a_{\mathbf{k}}^* a_{\mathbf{k}}\rangle + 3/2 \qquad (3)$$

if we choose $\varepsilon = 8\alpha/3\pi K$, which is smaller and better by a factor 1/3 from the ε in the article. Here we have employed an orthogonal rotation of coordinates bringing $\sum_{|\mathbf{k}|\geq K} a_{\mathbf{k}}^* a_{\mathbf{k}}$ into a form $(4/\varepsilon)\mathbf{Z}^*\mathbf{Z}$+non-negative operators. (Compare Eqs.(21,22) of the article.) Combining these inequalities, we obtain

$$-(\frac{4\pi\alpha}{V})^{1/2}\sum_{|\mathbf{k}|\geq K}\left[\langle \frac{a_{\mathbf{k}}}{|\mathbf{k}|}e^{i\mathbf{k}\cdot\mathbf{x}}\rangle + c.c.\right] \leq \varepsilon\langle \mathbf{p}^2\rangle + \sum_{|\mathbf{k}|\geq K}\langle a_{\mathbf{k}}^* a_{\mathbf{k}}\rangle + 3/2. \qquad (4)$$

This last inequality is our replacement for the ultraviolet bound (10). It follows that $H \geq H_K - 3/2$ where H_K is as in Eq.(11), but with the coefficient of $\mathbf{p}^2$ given by $(1-8\alpha/3\pi K)$ rather than $(1-8\alpha/\pi K)$. With the choice $K = 8\alpha/3\pi$, inequality (13) becomes $H \geq -(16\alpha^2/3\pi^2) - 3/2$, a bound at least consistent with a known *upper* bound for the ground state energy linear in α.

The remainder of the article is an analysis of H_K and needs only minor modification. The coefficient of $\mathbf{p}^2$ in Eqs.(19,23,27,28,30) should be $(1 - c_1\alpha^{-1/5}/3)$ and, at the end of the article, $c_5 = (c_1/3 + 2c_4)c_P$. Due to the smaller value of ε defined above, our estimate on the coefficient of $\alpha^{9/5}$ in (31) is slightly improved to 2.337, rather than 3.822 as reported. Of course, our lower bound for the ground state energy is decreased merely by the constant $-3/2$, which is unimportant on a scale of $\alpha^{9/5}$.