# On Spectral Properties of Periodic Polyharmonic Matrix Operators.

Yu. E. Karpeshina *

ABSTRACT We consider a matrix operator $H = (-\Delta)^l + V$ in $R^n$, where $n \geq 2$, $l \geq 1$, $4l > n+1$, and $V$ is the operator of multiplication by a periodic in $x$ matrix $V(x)$. We study spectral properties of $H$ in the high energy region. Asymptotic formulae for Bloch eigenvalues and the corresponding spectral projections are constructed. The Bethe-Sommerfeld conjecture, stating that the spectrum of $H$ can have only a finite number of gaps, is proved.

## 1 Introduction.

We consider the operator

$$H = (-\Delta)^l + V \tag{1}$$

in $L_2(R^n)^m$, where $m \geq 1$, $n \geq 2$, $l \geq 1$, $4l > n+1$, $(-\Delta)^l$ is a diagonal $m \times m$ matrix, its diagonal elements being the scalar polyharmonic operators, and $V$ is the operator of multiplication by periodic in $x$ matrix $V(x)$. For the sake of simplicity, we assume that $V(x)$ has orthogonal periods $a_1, ..., a_n$, however all the results are also valid for nonorthogonal periods. Without loss of generality we consider that

$$V(x) = V_0 + \alpha \tilde{V}(x),$$

where $V_0$ is a diagonal matrix with entries not depending on $x$:

$$(V_0)_{qs} = v_{0q}\delta_{qs}, \quad q, s = 1, ..., m,$$

$$v_{01} \leq v_{02} \leq ... \leq v_{om};$$

a parameter $\alpha$ belongs to $[0, 1]$, and $\tilde{V}(x)$ satisfies the condition:

$$\int_Q \tilde{V}(x)dx = 0,$$

*Research partially supported by USNSF Grant DMS-9803498.

$Q$ being the elementary cell of the periods: $Q = [0, a_1] \times ... \times [0, a_n]$. We assume that $V(x)$ is infinitely differentiable.

The scalar case of $(-\Delta)^l + V$, $4l > n+1$, was studied in [Sk1, Sk2, K1, K2]. The Bethe-Sommerfeld conjecture, stating that the spectrum of $H$ can have only a finite number of gaps, was proved in [Sk1, Sk2] for the case of a rational lattice and a bounded potential. It was proved in [K1, K2] for the case of a general lattice and a general class of potentials. The case of the Schrödinger operator $l = 1$, $n \geq 3$ is not considered here. The reader can find the proof of the conjecture for the scalar case $n = 3$, $l = 1$ in [Sk3, V, K3, K2, HM], and for the more general situation $n \leq 4$ in [HM]. Recently, the results for the scalar cases $6l > n + 2$ [PS1, PS2] and $8l > d + 3$ [PS3] have appeared.

The aim of this paper is to construct perturbation formulae for eigenvalues and spectral projections in the high energy region and to prove the Bethe-Sommerfeld conjecture for the matrix operator (1). In the case, when the entries of $V_0$ are different from each other, the whole construction is analogous to the scalar case, see [K1, K2]. The scheme in [K1, K2] is based on the fact that the Bloch eigenvalues of the operator $(-\Delta)^l$ in $L_2(R^n)$ are not degenerated for almost all values of quasimomentum. In the matrix case, when $v_{os} = v_{oq}$ for at least one pair of $q, s, q \neq s$, the scalar scheme does not work, for the reason that the Bloch eigenvalues of the "unperturbed" operator $H_0 = (-\Delta)^l + V_0$ are degenerated for all values of quasimomentum. Here we will show how this case can be treated.

It is well known (see f.e. [G, ReSi]) that the spectral analysis of $H$ can be reduced to the study of a family of operators $H(t)$, $t \in K$, where $K$ is the elementary cell of the dual lattice,

$$K = [0, 2\pi a_1^{-1}) \times [0, 2\pi a_2^{-1}) \times ... \times [0, 2\pi a_n^{-1}).$$

Vector $t$ is called a *quasimomentum*. An operator $H(t)$, $t \in K$, acts in $L_2(Q)^m$. It is described by the formula (1) and the quasiperiodic conditions:

$$\vec{u}(x_1, ..., x_{r-1}, a_r, x_{r+1}...x_n) = \exp(it_r a_r)\vec{u}(x_1, ..., x_{r-1}, 0, x_{r+1}, ..., x_n), \qquad (2)$$

$$r = 1, 2, ..., n.$$

The derivatives with respect to $x_r$ must also satisfy the analogous conditions. The operator $H(t)$, $t \in K$, has a discrete semibounded below spectrum $\Lambda(t)$:

$$\Lambda(t) = \cup_{s=1}^{\infty} \lambda_s(t), \ \lambda_s(t) \to_{s\to\infty} \infty.$$

The spectrum $\Lambda$ of the operator $H$ is the union of the spectra of the operators $H(t)$: $\Lambda = \cup_{t\in K}\Lambda(t) = \cup_{s\in N, t\in K}\lambda_s(t)$. The functions $\lambda_s(t)$ are continuous, so $\Lambda$ has a band structure:

$$\Lambda = \cup_{s=1}^{\infty}[q_s, Q_s], \ q_s = \min_{t\in K} \lambda_s(t), \ Q_s = \max_{t\in K} \lambda_s(t).$$

Absolute continuity of the spectrum was proven in [Th] for the case of the scalar Schrödinger operator and the matrix case recently in [KuLe]).

The eigenfunctions of $H(t)$ and $H$ are simply related. Extending all the eigenfunctions of the operators $H(t)$ quasiperiodically (see (2)) to $R^n$, we obtain a complete system of eigenfunctions of $H$.

We use the following representation of the potential:

$$\tilde{V}_{qs}(x) = \sum_{j \in Z^n, j \neq 0} v_{qs,j} \exp i(\vec{p}_j(0), x), \quad q, s = 1, ..., m, \tag{3}$$

where $\tilde{V}_{qs}(x)$ are matrix elements of $\tilde{V}(x)$, $v_{qs,j}$, $j \in Z^n$, are Fourier coefficients of $\tilde{V}_{qs}(x)$, $(\cdot, \cdot)$ is the inner product in $R^n$, $\vec{p}_j(0)$ are vectors of the dual lattice:

$$\vec{p}_j(0) = 2\pi(j_1 a_1^{-1}, j_2 a_2^{-1}, ..., j_n a_n^{-1}), \ j \in Z^n. \tag{4}$$

In the case $\tilde{V} = 0$, the eigenvalues and eigenfunctions of the operators $H_0(t)$, $t \in K$, are naturally indexed by pairs of indices $(j, q)$, $j \in Z^n$, $q = 1, ..., m$:

$$\lambda^0_{j,q}(t) = p_j^{2l}(t) + v_{0q}, \quad \psi^0_{j,q}(t, x) = \frac{1}{|Q|^{1/2}} \vec{e}_q^{\,0} \exp i(\vec{p}_j(t), x),$$

here $\vec{e}_q^{\,0}$ are vectors of the canonical basis in $R^m$: $(\vec{e}_q^{\,0})_s = \delta_{qs}, s = 1, ..., m$, and $\vec{p}_j(t) = \vec{p}_j(0) + t$, $\vec{p}_j(0)$ being given by (4); $p_j^{2l}(t) = |\vec{p}_j(t)|^{2l}$, and the symbol $|Q|$ stands for the volume of $Q$.

The operator $H(t)$ admits the matrix representation in the basis $\left\{\psi^0_{j,q}\right\}_{j \in Z^n, q=1,...,m}$:

$$H_{(j,q)(i,s)} = p_j^{2l}(t)\delta_{ji}\delta_{qs} + v_{oq}\delta_{ji}\delta_{qs} + v_{qs,j-i}.$$

In physical literature, the important concept of an isoenergetic surface of a periodic operator is used (see f.e. [Ki, Ma]). It is said that a point $t$ belongs to the isoenergetic surface $S(k)$ of the operator $H$, if the operator $H(t)$ has an eigenvalue equal to $k^{2l}$, i.e., if there is an $s$ such that $\lambda_s(t) = k^{2l}$. Let us consider the isoenergetic surface of the operator $H_0 = (-\Delta)^l + V_0$. The isoenergetic surface $S_0(k)$ of $H_0$ is the set in $K$, such that every $t \in S_0(k)$ satisfies the equation $p_j^{2l}(t) + v_{oq} = k^{2l}$ at least for one pair $(j, q)$. The surface can be obtained as follows: the spheres of the radiuses $\left(k^{2l} - v_{oq}\right)^{\frac{1}{2l}}, q = 1, ..., m$ centered at the origin of $R^n$, are divided into pieces by the dual lattice $\{\vec{p}_j(0)\}_{j \in Z^n}$, and, then, all these pieces are shifted into the elementary cell $K$ of the dual lattice. Thus, we obtain $m$ spheres "packed into the bag" $K$. If $v_{oq} = v_{os}$ for $q \neq s$, then the corresponding spheres coincide. Thus, the real number of spheres is equal to the number of <u>different</u> entries in $V_0$. Each point of a degenerated sphere corresponds to a multiple eigenvalue. The selfintersections of $S_0(k)$ are described by the equations: $p_j^{2l}(t) + v_{oq} = p_i^{2l}(t) + v_{os}$, $j, i \in Z^n, j \neq i$, $q, s = 1, ..., m$.

Let $S_1(k) \subseteq S_0(k)$. We say that $S_1(k)$ has an asymptotically full measure on $S_0(k)$ if the relation

$$\frac{s(S_1(k))}{s(S_0(k))} \to_{k \to \infty} 1 \tag{5}$$

holds, where $s(\cdot)$ is the area of a set, counting the multiplicity of the spheres.

## 2 The main results.

For any $t \in S_0(k)$ there is a pair $(j,q)$ such that

$$p_j^{2l}(t) + v_{oq} = k^{2l}, \tag{6}$$

i.e., $H_0(t)$ has an eigenvalue equal $k^{2l}$. The multiplicity of this eigenvalue is at least equal to the multiplicity $m_q$ of $v_{0q}$ as an entry in $V_0$. Let us consider the situation when it is greater. In fact, all other eigenvalues of $H_0$ can be represented as either $p_j^{2l}(t) + v_{os}$, $v_{os} \neq v_{oq}$, or $p_i^{2l}(t) + v_{os}, i \neq j$, $s$ being any number of $1, ..., m$ including $q$. The eigenvalues of the first type are separated from (6) by the distance not less than

$$d_0 = \min_{s,r:v_{os} \neq v_{or}} |v_{os} - v_{or}|. \tag{7}$$

An eigenvalue of the second type may coincide with (6). In other words, (6) has the multiplicity greater than $m_s$ if and only if there are $i \neq j$ and $s = 1, ..., m$, such that

$$p_j^{2l}(t) + v_{oq} = p_i^{2l}(t) + v_{os}. \tag{8}$$

This condition describes a selfintersection of the isoenergetic surface $S_0(k)$. Obviously, the set of the points $t$ satisfying (8) has the measure zero on $S_0(k)$.

We introduce a set $\chi_0(k,\delta)$, $\delta$ being a positive parameter, as $S_0(k) \setminus A_0(k,\delta)$, where $A_0(k,\delta)$ is the $(k^{-n+1-\delta})$-neighborhood of all the selfintersections (8) of $S_0(k)$. The properties of $\chi_0(k,\delta)$ are described by the following lemma.

**Lemma 1** *If $t$ belongs to $\chi_0(k,\delta)$, then there are a unique index $j$ and a unique value $v_{0q}$, such that (6) holds. Moreover, for all $i \neq j$ and $s = 1, ..., m$:*

$$\left| p_j^{2l}(t) + v_{0q} - p_i^{2l}(t) - v_{0s} \right| > 2k^{2l-n-\delta}. \tag{9}$$

*The set $\chi_0(k,\delta)$ has an asymptotically full measure on $S_0(k)$.*

Lemma 1 is stable in the $(k^{-n+1-\mathbf{2}\delta})$-neighborhood of of $\chi_0(k,\delta)$ in the sense that, those points, which are close to $\chi_0(k,\delta)$, satisfy the equation (6) approximately and also satisfy (9). The following generalization of Lemma 1 holds:

**Lemma 2** *If $t$ belongs to the $(k^{-n+1-\mathbf{2}\delta})$-neighborhood of of $\chi_0(k,\delta)$, then there are a unique index $j$ and a unique value $v_{0q}$, such that*

$$|p_j^{2l}(t) + v_{0q} - k^{2l}| < 3lk^{2l-n-\mathbf{2}\delta} \tag{10}$$

*and the inequality (9) holds for all $i \neq j$ and $s = 1, ..., m$. The set $\chi_0(k,\delta)$ has an asymptotically full measure on $S_0(k)$.*

Thus, if $t \in \chi_0(k,\delta)$, then there is only one index $j = j(t)$ and there is only one value $v_{oq}$ satisfying (6). When all entries of $V_0$ are different from each other, the index $q = q(t)$ is defined uniquely by $v_{0q}$. If there are multiple entries in $V_0$, we define $q(t)$ as the smallest of all indices satisfying (6). By this rule, $q(t)$ is defined uniquely. When $t$ is in the $(k^{-n+1-2\delta})$-neighborhood of $\chi_0(k,\delta)$, the indices $j(t)$ and $q(t)$ are uniquely defined by inequality (10). We denote by $m_q$ the multiplicity of $v_{0q}$, i.e., $m_q : v_{0,q-1} < v_{0,q} = .... = v_{0,q+m_q-1} < v_{0,q+m_q}$.

**Corollary 1** *If $2l \leq n$, and $t$ is in the $(k^{-n+1-2\delta})$-neighborhood of $\chi_0(k,\delta)$, then the operator $H_0(t)$ has a unique eigenvalue satisfying (10). It has the multiplicity $m_q$. The distance from $p_j^{2l}(t) + v_{0q}$ to the nearest other eigenvalue of $H_0(t)$ is not less than $2k^{2l-n-\delta}$, for $k$ large enough.*

In fact, according to the lemma, the eigenvalues of the second type are separated from $p_j^{2l}(t)+v_{0q}$ at least by the distance $2k^{2l-n-\delta}$. The eigenvalues of the first type are separated at least by the distance $d_0$ (see (7)), which does not depend on $k$ and obviously greater than $2k^{2l-n-\delta}$ for $k$ large enough.

**Corollary 2** *If $2l > n$, $0 < \delta < 2l-n$, and $t$ is in the $(k^{-n+1-2\delta})$-neighborhood of $\chi_0(k,\delta)$, then the operator $H_0(t)$ has a unique eigenvalue satisfying (10). It has the multiplicity $m_q$. The distance from $p_j^{2l}(t)+v_{0q} = k^{2l}$ to the nearest other eigenvalue of $H_0(t)$ is not less than $d_0$, for $k$ large enough.*

This corollary is based on the fact that $d_0 < 2k^{2l-n-\delta}$ for $2l > n$, $0 < \delta < 2l-n$, and $k$ large enough.

Let us consider the circle $C_0$ in the complex plane, which is centered at $z = k^{2l}$ and has the radius $k^{2l-n-\delta}$:

$$C_0 = \{z : |z - k^{2l}| = k^{2l-n-\delta}\}.$$

If $2l \leq n$ and $t$ belongs to the $(k^{-n+1-2\delta})$-neighborhood of $\chi_0(k,\delta)$, then according to Corollary 1, the circle includes only one eigenvalue of $H_0(t)$. It has the multiplicity $m_q$. If $2l > n$, $0 < \delta < 2l-n$, then the circle includes all the eigenvalues given by the formula

$$p_j^{2l}(t) + v_{0s},\ s = 1, 2, ...m,$$

since $|v_{0s} - v_{0q}| < k^{2l-n-\delta}$. All other eigenvalues are outside the circle. Counting multiplicity, we have exactly $m$ eigenvalues of $H_0(t)$ inside the circle.

In the scalar situation $m = m_q = 1$, i.e., the circle has one simple eigenvalue of $H_0(t)$ inside. In the matrix situation we can have multiple eigenvalues, which makes it more difficult to describe their shift under the perturbation $\alpha\tilde{V}$.

If we can prove the convergence of a perturbation series for the resolvent of $H(t)$ on $C_0$, then the number of eigenvalues inside $C_0$ does not change with perturbation. The circle $C_0$ is chosen in such a way, that its distance from the eigenvalues of $H_0(t)$ is equal to $k^{2l-n-\delta}$, if $t \in \chi(k,\delta)$. When $2l > n$, $0 < \delta < 2l-n$, and $t$ is in the $(k^{-n+1-2\delta})$- neighborhood

of $\chi_0(k,\delta)$, then the perturbation series converges on $C_0$, since the distance from $C_0$ to the unperturbed eigenvalues is greater than $\|\tilde{V}\|$ for $k$ large enough. When $2l \le n$ then, generally speaking, the perturbation series diverges for $t \in \chi_0(k,\delta)$, since the distance from $C_0$ to the eigenvalues is small (it goes to zero as $k \to \infty$). We construct a "nonsingular" set $\chi_1 \subset \chi_0(k,\delta)$, such that it provides the convergence of the series in the case $2l \le n$ and also improves the convergence in the case $2l > n$.

**Lemma 3** *Let $0 < \beta < 1$, and $0 < 2\delta < (n-1)(1-\beta)$. Then, for any sufficiently large $k$, $k > k_0(\beta,\delta,a_1,...a_n)$, there exists a set $\chi_1(k,\beta,\delta) \subset \chi_0(k,\delta) \subset S_0(k)$, such that for any point $t$ in the $(k^{-n+1-2\delta})$-neighborhood of this set the following inequality holds:*

$$\min_{z\in C_0} \min_{\substack{i,i' \in Z^n, 0<|i-i'|<k^\beta, \\ s,s'=1,...m}} |\, p_i^{2l}(t) + v_{0s} - z \,||\, p_{i'}^{2l}(t) + v_{0s'} - z \,| > k^{2\gamma}, \tag{11}$$

$$2\gamma = 4l - n - 1 - \beta(n-1) - 2\delta.$$

*The set $\chi_1(k,\beta,\delta)$ has an asymptotically full measure on $S_0(k)$. Moreover,*

$$\frac{s(S_0(k) \setminus \chi_1(k,\beta,\delta))}{s(S_0(k))} =_{k\to\infty} O(k^{-\delta/8}). \tag{12}$$

Further we assume: $0 < \beta < \beta_0$, $\beta_0 = \min\{1, (4l-n-1)(n-1)^{-1}\}$, and $0 < \delta < \delta_0$, $\delta_0 = \frac{1}{4}\min\{(n-1)(1-\beta), 4l-n-1-\beta(n-1)\}$. Obviously, under these conditions $\gamma$ is positive.

The set $\chi_1(k,\beta,\delta)$ provides the convergence of the series for the resolvent:

**Theorem 1** *If $t$ belongs to the $(k^{-n+1-2\delta})$-neighborhood of $\chi_1(k,\beta,\delta)$ in $K$, then the perturbation series for the resolvent*

$$(H(t)-z)^{-1} = (H_0(t)-z)^{-1} \sum_{r=0}^{\infty} \left(-\alpha \tilde{V}(H_0(t)-z)^{-1}\right)^r \tag{13}$$

*converges in the class of bounded operators for any $z \in C_0$.*

This theorem enable us to obtain asymptotic formulae for Bloch eigenvalues and the corresponding spectral projections. The next two theorems provide information about the location of eigenvalues.

**Theorem 2** *If $2l \le n$ and $t$ belongs to the $(k^{-n+1-2\delta})$-neighborhood of $\chi_1(k,\beta,\delta)$, then there are exactly $m_q$, $q = q(t)$, eigenvalues of the operator $H(t)$ in the interval $\Delta_0 = [p_j^{2l}(t) + v_{0q} - k^{2l-n-\delta}, p_j^{2l}(t) + v_{0q} + k^{2l-n-\delta}]$. In fact, they are located in the smaller interval $\Delta_1 = [p_j^{2l}(t) + v_{0q} - k^{-2l+1+\zeta}, p_j^{2l}(t) + v_{0q} + k^{-2l+1+\zeta}]$, $\zeta = \beta(n-1) + 2\delta$. The corresponding spectral projection is given by the series:*

$$E(t) = E_0 + \sum_{r=1}^{\infty} \alpha^r G_r(k,t), \tag{14}$$

*where $E_0$ is the unperturbed spectral projection:*

$$E_0 = \sum_{s=q}^{q+m_q-1} E_{0s}, \quad (E_{0s})_{(ip)(i'p')} = \delta_{ii'}\delta_{ji}\delta_{pp'}\delta_{ps}, \tag{15}$$

*the operators $G_r$ are given by the formula*

$$G_r(k,t) = \frac{(-1)^{r+1}}{2\pi i}\oint_{C_0}((H_0(t)-z)^{-1}\tilde{V})^r(H_0(t)-z)^{-1}dz, \quad r=1,2..., \tag{16}$$

*and satisfy the estimates:*

$$\| G_r(k,t) \| < c_V^r k^{-\gamma r}, \quad r=1,2..., \tag{17}$$

*$c_V$ not depending on $k$, $\|\cdot\|$ being the norm in the class of bounded operators.*

**Theorem 3** *If $2l > n$ and $t$ belongs to the $(k^{-n+1-2\delta})$-neighborhood of $\chi_1(k,\beta,\delta)$, then there are exactly $m$ eigenvalues of the operator $H(t)$ in the interval $\Delta_0 = [p_j^{2l}(t) + v_{0q} - k^{2l-n-\delta}, p_j^{2l}(t) + v_{0q} + k^{2l-n-\delta}]$. In fact, they are located in the smaller intervals $\Delta_s = [p_j^{2l} + v_{0s} - k^{-2l+1+\zeta}, p_j^{2l} + v_{0s} + k^{-2l+1+\zeta}]$, $\zeta = \beta(n-1)+2\delta$, $s = 1,...m$. The corresponding spectral projection is given by the series*

$$E(t) = E_0 + \sum_{r=1}^{\infty}\alpha^r G_r(k,t), \tag{18}$$

*where $E_0$ is the unperturbed spectral projection:*

$$E_0 = \sum_{s=1}^{m} E_{0s}, \quad (E_{0s})_{(ip)(i'p')} = \delta_{ii'}\delta_{ji}\delta_{pp'}\delta_{ps}, \tag{19}$$

*the operators $G_r$ are given by formula (16) and satisfy the estimates (17).*

*Remark.* It is possible to construct the perturbation series for each interval $\Delta_s$ separately, however it gives slower convergence.

The proof of the Bethe-Sommerfeld conjecture for the scalar polyharmonic operator in [K1, K2] is essentially based on the fact that there is a unique simple eigenvalue in the interval $\Delta_0$. Under such circumstances it is possible to differentiate the eigenvalue with respect to $t$, to obtain a suitable estimate from below for the derivative and to prove, using the estimate, that any sufficiently large $k^{2l}$ belongs to the spectrum of $H$. In the present case we do not have, generally speaking, the uniqueness of eigenvalue. However, it is still possible to get the estimates for the derivatives of eigenvalues with respect to $t$ and to prove the following theorem.

**Theorem 4** *(The proof of the Bethe-Sommerfeld conjecture) There is only a finite number of gaps in the spectrum of the operator $H$.*

# 3 The proofs.

The proof of Lemma 2, up to minor technical details, coincides with the proof of Lemma 2.5 in [K2]. For the scalar case the set $\chi_1(k,\beta,\delta)$ is explicitly constructed in [K2] (Lemma 3.3) and it is proved there (Lemma 3.6) that $\chi_1(k,\beta,\delta)$ has an asymptotically full measure in the sense (12). The proof of Lemma 2 is analogous to the proof in [K2].

The proof of Theorem 1. Let us rewrite the series (13) in the form:

$$(H(t)-z)^{-1} = (H_0(t)-z)^{-1} + \sum_{r=1}^{\infty} (H_0(t)-z)^{-1/2}(-\alpha A)^r (H_0(t)-z)^{-1/2}, \qquad (20)$$

$$A = (H_0(t)-z)^{-1/2}\tilde{V}(H_0(t)-z)^{-1/2}.$$

We prove that the perturbation series converges for any $t$, satisfying (9) and (11). Inequality (9) and the definition of $C_0$ yields the estimate:

$$\|(H_0-z)^{-1/2}\| = \left(\min_{i\in Z^n,\ s=1,2,\dots m} |p_i^{2l}(t)+v_{0s}-z|\right)^{-1/2} = k^{-(2l-n-\delta)/2}. \qquad (21)$$

To prove the convergence of the series for a large $k$, it is enough to check that

$$\|A\| < c_V k^{-\gamma},\ z\in C_0, \qquad (22)$$

$C_V$ not depending on $k$. The matrix elements of the operator $A$ are given by the formula:

$$A_{(is)(i's')} = \frac{v_{ss',i-i'}}{\sqrt{\left(p_i^{2l}(t)+v_{0s}-z\right)\left(p_{i'}^{2l}(t)+v_{0s'}-z\right)}}, \qquad (23)$$

if $i\neq i'$ and are equal to zero if $i=i'$. We represent $A$ in the form $A=A'+A''$, where

$$A'_{(is)(i's')} = \begin{cases} A_{(is)(i's')}, & \text{if } 0<|i-i'|<k^\beta, \\ 0, & \text{otherwise,} \end{cases}$$

and, correspondingly,

$$A''_{(is)(i's')} = \begin{cases} 0, & \text{if } |i-i'|<k^\beta, \\ A_{(is)(i's')}, & \text{otherwise.} \end{cases}$$

Considering the estimate (11), we see that the denominator on the right-hand side of (23) can be bounded from below by $k^\gamma$, when $0<|i-i'|<k^\beta$. Therefore,

$$\left|A'_{(is)(i's')}\right| < |v_{ss',i-i'}|k^{-\gamma}.$$

Taking into account that $v_{ss',i-i'}$ are the Fourier coefficients of infinitely differentiable functions, we obtain

$$\sup_{s,s'=1,\dots m}\ \sum_{i\in Z^n,|i-i'|<k^\beta} |v_{ss',i-i'}| < \tilde{c}_V,$$

and, therefore,

$$\|A'(z)\| < c_V k^{-\gamma},\ z \in C_0. \tag{24}$$

Using (21), we obtain:

$$\left|A''_{(is)(i's')}\right| < |v_{ss',i-i'}|k^{-(2l-n-\delta)}.$$

Considering that $V$ is infinitely differentiable, we obtain:

$$\sup_{s,s'=1,\dots m} \sum_{i\in Z^n, |i-i'|>k^\beta} |v_{ss',i-i'}| < c_V k^{2l-n-\delta-\gamma},\quad c_V = c_V(\delta,\beta,\gamma),$$

and, therefore,

$$\|A''(z)\| < c_V k^{-\gamma},\ z \in C_0. \tag{25}$$

Adding (24) and (25), we obtain (22). The theorem is proved.

The proof of Theorem 2. The operator $H_0$ has $m_q$ eigenvalues inside $C_0$. The fact that $(H(t)-z)^{-1}$ is bounded on $C_0$ for all $\alpha$ means that the number of eigenvalues inside $C_0$ does not change with the perturbation $\alpha\tilde{V}$, $\alpha \in [0,1]$. Since all eigenvalues are real, they are inside the interval $\Delta_0$, which is cut out by the circle on the real axis. Integrating the resolvent over the contour, we get the corresponding spectral projection:

$$E(t) = -\frac{1}{2\pi i}\oint_{C_0}(H(t)-z)^{-1}dz. \tag{26}$$

Substituting the series for the resolvent (13) in the last formula, we obtain (14)–(16). Considering the estimates (21) and (22), we get (17).

We prove that, in fact, the perturbation series for the resolvent converges on a smaller circle $C_1$, which has the radius $k^{-2l+1+\zeta}$ and the same center $z = k^{2l}$ as $C_0$. Let $z \in C_1$ and $z_0 \in C_0$. For all $i \neq j$:

$$|p_i^{2l}(t) + v_{0s} - z| > |p_i^{2l}(t) + v_{0s} - z_0|, \tag{27}$$

since the circle $C_1$ is smaller than $C_0$ and, therefore, further away from outside eigenvalues. Since (11) holds for $z_0$, we easily obtain:

$$|p_i^{2l}(t) + v_{0s} - z||p_{i'}^{2l}(t) + v_{0s'} - z| > k^{2\gamma} \quad \text{when} \quad i,i' \neq j,\ 0 < |i-i'| < k^\beta.$$

Suppose $i = j, i' \neq j$. Then, using (27) for $i'$, we get:

$$|p_j^{2l}(t) + v_{0s} - z||p_{i'}^{2l}(t) + v_{0s'} - z| >$$

$$\frac{|p_j^{2l}(t) + v_{0s} - z|}{|p_j^{2l}(t) + v_{0s} - z_0|}|p_j^{2l}(t) + v_{0s} - z_0||p_{i'}^{2l}(t) + v_{0s'} - z_0|.$$

Using the definitions of $C_0$ and $C_1$, we obtain:

$$\frac{|p_j^{2l}(t) + v_{0s} - z|}{|p_j^{2l}(t) + v_{0s} - z_0|} = \frac{k^{-2l+1+\zeta}}{k^{2l-n-\delta}} = k^{-2\gamma+\delta}.$$

Taking into account (11) for $z_0$, we arrive at the inequality:

$$|p_j^{2l}(t) + v_{0s} - z||p_{i'}^{2l}(t) + v_{0s'} - z| > k^{\delta}.$$

Thus, for any $z \in C_1$

$$\min_{\substack{i,i' \in Z^n, 0 < |i-i'| < k^{\beta}, \\ s,s' = 1,...m}} | p_i^{2l}(t) + v_{0s} - z \,||\, p_{i'}^{2l}(t) + v_{0s'} - z \,| > k^{\delta}. \tag{28}$$

Considering as above, we prove that the perturbation series for the resolvent converges on the circle $C_1$ for all $\alpha \in [0,1]$. This means that all eigenvalues are, in fact, inside the interval $\Delta_1$, which is cut out by $C_1$ on the real axis. Note that the series for the resolvent and, therefore, for the spectral projection converges slower on $C_1$ than on $C_0$, since the estimate (22) valid on $C_0$ is replaced by a weaker estimate $\|A(z)\| < c_V k^{-\delta}$ on $C_1$. The theorem is proved.

The prove of Theorem 3 is analogous to that of Theorem 2. The only difference that we replace the circle $C_0$ by smaller circles $C_s$ which cut out the intervals $\Delta_s$ from the real axis.

Let $t_0 \in \chi_1(k,\beta,\delta)$ and $t = t_0 + \tau\vec{\nu}$, where $|\tau| < k^{-n+1-2\delta}$, and $\vec{\nu}$ is a unit vector in $R^n$. The eigenvalues $\lambda_s$, lying inside $C_0$, analytically depend on $\tau$. Let us estimate their derivatives.

**Lemma 4** *If $t_0 \in \chi_1(k,\beta,\delta)$ and $t = t_0 + \tau\vec{\nu}$, $|\tau| < k^{-n+1-2\delta}$, $\vec{\nu}$ being a unit vector in $R^n$, then the derivative of any $\lambda_s(\tau)$ with respect to $\tau$ satisfies the asymptotic estimate:*

$$\frac{\partial \lambda_s(\tau)}{\partial \tau} = 2p_j^{2l-2}(t)(\vec{p}_j(t),\vec{\nu}) + O(k^{2l-1-\gamma}), \tag{29}$$

*when $k \to \infty$.*

Proof. Let us assume for definiteness that $2l \le n$. There are exactly $m_q$ eigenvalues inside $\Delta_0$, counting the multiplicity. Let $\phi_s(\tau)$ be a normalized eigenvector corresponding to a $\lambda_s(\tau)$. [1] Then,

$$\frac{\partial \lambda_s}{\partial \tau} =< \frac{\partial H_0}{\partial \tau}\phi_s(\tau), \phi_s(\tau) >, \tag{30}$$

where $< \cdot, \cdot >$ is the inner product in $l_2(Z^n)^m$ and $\frac{\partial H_0}{\partial \tau}$ is a diagonal matrix:

$$\left(\frac{\partial H_0}{\partial \tau}\right)_{(ir)(i'r')} = 2p_i^{2l-2}(t)(\vec{p}_i(t),\vec{\nu})\delta_{ii'}\delta_{rr'}, \quad i,i' \in Z^n,\ r,r' = 1,...,m. \tag{31}$$

[1] An eigenvector $\phi_s(\tau)$ is defined up to $\pm$ sign if $\lambda_s(\tau)$ is simple. It is allowed to be any normalized eigenvector if the multiplicity of $\lambda_s(\tau)$ is greater than 1 for all $\tau$. We exclude from the consideration the points $\tau$, where two or more nonidentical eigenvalues coincide, since there is only a finite number of such points.

We denote by $\vec{e}^{\,0}_{jr}$ $r = q, ..., q + m_q - 1$ the eigenvectors of $H_0(t)$, corresponding to $p_j^{2l} + v_{0q}$: $(\vec{e}^{\,0}_{jr})_{(ir')} = \delta_{ij}\delta_{r'r}$ $i \in Z^n, r' = 1, ...m$. They form a basis in $E_0 l_2(Z^n)^m$. Let us consider the vectors $\vec{e}_{jr}$, $r = q, ..., q + m_q - 1$, defined by the formula: $\vec{e}_{jr} = E(t)\vec{e}^{\,0}_{jr}$. They form a complete almost orthogonal system in $E(t)l_2(Z^n)^m$, since

$$E(t) = E_0 + O(k^{-\gamma}) \tag{32}$$

(in the sense of bounded operators) and $E_0\vec{e}^{\,0}_{jr} = \vec{e}^{\,0}_{jr}$, and, therefore,

$$\vec{e}_{jr} = \vec{e}^{\,0}_{jr} + O(k^{-\gamma}). \tag{33}$$

Hence, an eigenvector $\phi_s(\tau)$ can be represented as a linear combination of $\vec{e}_{jr}$:

$$\phi_s(\tau) = \sum_{r=q}^{q+m_q-1} c_r^{(s)}\vec{e}_{jr}.$$

Considering that function $\phi_s(\tau)$ is normalized, we obtain

$$\sum_{r,r'=q}^{q+m_q-1} c_r^{(s)}\bar{c}_{r'}^{(s)} < \vec{e}_{jr}, \vec{e}_{jr'} >= 1,$$

and, by (33),

$$\sum_{r=q}^{q+m_q-1} |c_r^{(s)}|^2 = 1 + O(k^{-\gamma}). \tag{34}$$

Using again (33), we obtain

$$\phi_s(\tau) = \phi_s^0(\tau) + \delta\phi_s(\tau), \tag{35}$$

where $\phi_s^0(\tau) = \sum_{r=q}^{q+m_q-1} c_r^{(s)}\vec{e}^{\,0}_{jr}$, and $\delta\phi_s(\tau)$ satisfies the estimate:

$$\|\delta\phi_s(\tau)\| < c_V k^{-\gamma}, \tag{36}$$

$\|\cdot\|$ being the norm in $l_2(Z^n)^m$. Substituting (35) in (30), we obtain

$$\frac{\partial\lambda_s(\tau)}{\partial\tau} = < \frac{\partial H_0}{\partial\tau}\phi_s^0(\tau), \phi_s^0(\tau) > + < \frac{\partial H_0}{\partial\tau}\phi_s^0(\tau), \delta\phi_s(\tau) > + \tag{37}$$

$$< \frac{\partial H_0}{\partial\tau}\delta\phi_s(\tau), \phi_s^0(\tau) > + < \frac{\partial H_0}{\partial\tau}\delta\phi_s(\tau), \delta\phi_s(\tau) > .$$

Obviously,

$$< \frac{\partial H_0}{\partial\tau}\phi_s(\tau)^0, \phi_s^0(\tau) >= \sum_{r,r'=q}^{q+m_q-1} c_r^{(s)}\bar{c}_{r'}^{(s)} < \frac{\partial H_0}{\partial\tau}\vec{e}^{\,0}_{jr}, \vec{e}^{\,0}_{jr'} > .$$

Substituting (31), we get:

$$< \frac{\partial H_0}{\partial\tau}\phi_s^0(\tau), \phi_s^0(\tau) >= 2p_j^{2l-2}(t)(\vec{p}_j(t), \vec{\nu}) \sum_{r=q}^{q+m_q-1} |c_r^{(s)}|^2.$$

Taking into account (34), we obtain:

$$< \frac{\partial H_0}{\partial \tau}\phi_s^0(\tau), \phi_s^0(\tau) >= 2p_j^{2l-2}(t)(\vec{p}_j(t), \vec{\nu}) \left(1 + O(k^{-\gamma})\right). \tag{38}$$

Let us estimate $\|\frac{\partial H_0}{\partial \tau}\delta\phi_s(\tau)\|$. Using (31), is easy to show that

$$\left\|\frac{\partial H_0}{\partial \tau}\delta\phi_s\right\|^2 = 4 \sum_{i\in Z^n, r=1,\dots m} p_i^{4l-4}(t)(\vec{p}_i(t), \vec{\nu})^2 |(\delta\phi_s(\tau))_{ir}|^2.$$

We estimate the terms of the series as follows:

$$p_i^{4l-4}(t)(\vec{p}_i(t), \vec{\nu})^2 |(\delta\phi_s)_{ir}|^2 \le X_{ir}Y_{ir},$$

$$X_{ir} = p_i^{4l-2}(t)|(\delta\phi_s)_{ir}|^{2-\frac{1}{l}},$$

$$Y_{ir} = |(\delta\phi_s)_{ir}|^{\frac{1}{l}}.$$

Using Hölder's inequality with $p = \frac{2l}{2l-1}$, $q = 2l$, we obtain:

$$\left\|\frac{\partial H_0}{\partial \tau}\delta\phi_s(\tau)\right\|^2 \le \|H_0\delta\phi_s(\tau)\|^{\frac{2l-1}{l}} \|\delta\phi_s(\tau)\|^{\frac{1}{l}}. \tag{39}$$

It is easy to obtain an estimate for $\|H_0\delta\phi_s(\tau)\|$. In fact, $\phi_s(\tau)$ satisfies the equation $H(t)\phi_s(\tau) = \lambda_s\phi_s(\tau)$ and $\phi_s^0(\tau)$ satisfies the equation $H_0(t)\phi_s^0(\tau) = (p_j^{2l}(t)+v_{0q})\phi_s^0(\tau)$.Therefore, the function $\delta\phi_s$ satisfies the equation

$$H_0(t)\delta\phi_s(\tau) + \tilde{V}\phi_s(\tau) = (p_j^{2l}(t) + v_{0q})\delta\phi_s(\tau) + (\lambda_s - p_j^{2l}(t) - v_{0q})\phi_s(\tau).$$

Hence,

$$H_0(t)\delta\phi_s(\tau) = (p_j^{2l}(t) + v_{0q})\delta\phi_s(\tau) + f, \quad \|f\| < C_V,$$

and $\|H_0\delta\phi_s(\tau)\| < k^{2l}\|\delta\phi_s(\tau)\| + C_V$. Noting that $2l - \gamma > 0$ and using (36), we get: $\|H_0\delta\phi_s(\tau)\| < C_V k^{2l-\gamma}$. Substituting the last estimate in (39) and using again (36), we obtain:

$$\left\|\frac{\partial H_0}{\partial \tau}\delta\phi_s(\tau)\right\| \le c_V k^{2l-1-\gamma}. \tag{40}$$

Using (36) and (40), we arrive at the inequality:

$$\left| < \frac{\partial H_0}{\partial \tau}\phi_s^0(\tau), \delta\phi_s(\tau) > + < \frac{\partial H_0}{\partial \tau}\delta\phi_s(\tau), \phi_s^0(\tau) > + \right.$$

$$\left. < \frac{\partial H_0}{\partial \tau}\delta\phi_s(\tau), \delta\phi_s(\tau) > \right| < c_V k^{2l-1-\gamma}. \tag{41}$$

Combining (30), (37), (38) and (41), we get (29). The analogous considerations give the proof of the lemma for the case $2l > n$. The lemma is proved.

**Corollary 3** *Suppose that $\vec{\nu}$ is chosen in such a way that $|(\vec{p}_j(t), \vec{\nu})| > p_j(t)/2$. Then,*

$$\frac{\partial \lambda_s}{\partial \tau} = 2p_j^{2l-2}(t)(\vec{p}_j(t), \vec{\nu}) \left(1 + O(k^{-\gamma})\right). \tag{42}$$

In fact, it is easy to see that the first term in the asymptotic formula (29) is the main one, since it satisfies the inequality $\left|2p_j^{2l-2}(t)(\vec{p}_j(t), \vec{\nu})\right| > k^{2l-1}$. Therefore, (42) holds.

*Remark.* The right part of (42) does not essentially depend on $s$. This means that, when $\tau$ changes, the eigenvalues $\lambda_s(\tau)$ lying in $\Delta_0$ move essentially as a group – all together.

The proof of Theorem 4. It is enough to prove that all $k^{2l}$, which are large enough, belong to the spectrum of $H$. Let $t_0 \in \chi_1(k, \beta, \delta)$. There is a pair $(j, q)$ such that $p_j^{2l}(t_0) + v_{0q} = k^{2l}$. Let $\vec{\nu}$ will be a unit vector in the direction of $\vec{p}_j(t_0)$. According to Theorems 2 and 3, for any $t = t_0 + \tau\nu$, $|\tau| < k^{-n+1-2\delta}$ and $k$ large enough, there are exactly $m_q$ eigenvalues of $H(t)$, which satisfy the estimate:

$$|\lambda_s(\tau) - (p_j^{2l}(t) + v_{0q})| < c_V k^{-2l+1+\zeta}. \tag{43}$$

The functions $\lambda_s(\tau)$ analytically depend on $\tau$. Using Corollary 3 and the special choice of vector $\vec{\nu}$, we obtain:

$$\frac{\partial \lambda_s(\tau)}{\partial \tau} = 2p_j^{2l-1}(t)\left(1 + o(1)\right), \tag{44}$$

for any eigenvalue $\lambda_s$ satisfying (43). We consider a $\lambda_s(\tau)$ as a function of $\tau$ in the domain $[-\tau_0, \tau_0]$, $\tau_0 = k^{-n+1-2\delta}$. It follows from (44) that $\lambda_s(\tau)$ is a monotonic function. Since it is continuous, its range is the interval $[\lambda_s(-\tau_0), \lambda_s(\tau_0)]$. Taking into account (43), we obtain:

$$\lambda_s(-\tau_0) = p_j^{2l}(t_0 - \tau_0\vec{\nu}) + v_{0q} + O(k^{-2l+1+\zeta}) = (k_0 - \tau_0)^{2l} + v_{0q} + O(k^{-2l+1+\zeta}),$$

$$k_0 = p_j(t_0) = \left(k^{2l} - v_{0q}\right)^{\frac{1}{2l}}.$$

Similarly,

$$\lambda_s(\tau_0) = p_j^{2l}(t_0 + \tau_0\vec{\nu}) + v_{0q} + O(k^{-2l+1+\zeta}) = (k_0 + \tau_0)^{2l} + v_{0q} + O(k^{-2l+1+\zeta}).$$

It is easy to observe that $k_0^{2l} + v_{0q} = k^{2l}$ and $k_0^{2l-1}\tau_0 = k^{2l-n-2\delta} > k^{-2l+1+\zeta}$. Therefore the interval $[\lambda_s(-\tau_0), \lambda_s(\tau_0)]$ contains $[k^{2l} - k^{2l-1}\tau_0, k^{2l} + k^{2l-1}\tau_0]$ when $k$ is large enough, and, obviously, contains $k^{2l}$. Thus, there is a value $\tau_s$ such that $\lambda_s(\tau_s) = k^{2l}$, i.e., the point $\lambda = k^{2l}$ belongs to the spectrum. The theorem is proved.

# References


[G] Gel'fand, I.M. *Expansion in Eigenfunctions of an Equation with Periodic Coefficients.* Dokl. Akad. Nauk SSSR, **73** (1950), pp. 1117-1120. (in Russian).

[HM] Helffer B., Mohamed A. *Asymptotics of the density of states for the Schrödinger operator with periodic electric potential.* Duke Math. J. **92** (1998), pp. 1 – 60.

[K1] Karpeshina, Yu. E. *Analytic Perturbation Theory for a Periodic Potential,* Izv. Akad. Nauk SSSR Ser. Mat., **53** (1989), 1, pp. 45-65; English transl.: Math. USSR Izv., **34** (1990), 1, pp. 43 – 63.

[K2] Karpeshina Yu.E. *Perturbation theory for the Schrödinger operator with a periodic potential*, in series "Lecture Notes in Mathematics", # 1663, Springer-Verlag, 1997, 352 pp.

[K3] Karpeshina, Yu.E. *Perturbation Theory for the Schrödinger Operator with a Periodic Potential.* Trudy Mat. Inst. Steklov, **188** (1990), pp. 88 – 116; Engl. transl.: Proceedings of the Steklov Institute of Mathematics, 1991, Issue 3, pp. 109-145.

[Ki] Kittel, Ch. *Introduction to Solid State Physics.* New-York, Wiley, c1976.

[KuLe] Kuchment, P., Levendorskii, S *On the structure of the spectra of periodic elliptic operator*

[Ma] Madelung, O., *Introduction to Solid State Theory.* Berlin, New-York, Springer-Verlag, 1978.

[PS1] Parnovski, L., Sobolev, A. V. *On the Bethe-Sommerfeld Conjecture.* quations aux Drives Partielles (La Chapelle sur Erdre, 2000), Exp. No. XVII, 13 pp., Univ. Nantes, Nantes, 2000.

[PS2] Parnovski, L., Sobolev, A.V. *On the Bethe-Sommerfeld Conjecture for the Polyharmonic Operator.* to appear in Duke Math. J.

[PS3] Parnovski, L., Sobolev, A.V., preprint.

[ReSi] Reed, M., Simon, B. *Methods of Modern Mathematical Physics.*, Vol IV, Academic Press, 3rd ed., New York – San Francisco – London (1987).

[Sk1] Skriganov, M.M. *Finiteness of the Number of Lacunae in the Spectrum of the Multidimensional Polyharmonic Operator with a Periodic Potential.* Mat. Sb., **113(155)** (1980), pp. 133-145; Engl. transl.: Math. USSR. Sb., **41** (1982), pp. 115 – 125.

[Sk2] Skriganov, M.M. *Geometric and Arithmetic Methods in the Spectral Theory of Multidimensional Periodic Operators.* Trudy Mat. Inst. Steklov., **171**, pp. 1 – 121. (1985); Engl. transl.: Proc. Steklov Inst. Math., **171**, 2 (1987).

[Sk3] Skriganov, M.M. *The Spectrum Band Structure of the Three-Dimensional Schrödinger Operator with a Periodic Potential.* Invent. Math., **80** (1985) pp. 107 –121.

[Th] Thomas, L.E. *Time Dependent Approach to Scattering from Impurities in Crystal.* Comm. Math. Phys., **33** (1973), pp. 335 – 343.

[V] Veliev, O.A. *Asymptotic Formulae for Eigenvalues of a Periodic Schrödinger Operator and Bethe-Sommerfeld Conjecture.* Functional. Anal. i Prilozhen., **21** (1987), no. 2, 1–15; Engl. transl.: Functional Anal. Appl., **21** (1987), pp. 87 –99.